

# *Positive Weighted Sectional Curvature*

LEE KENNARD & WILLIAM WYLIE

ABSTRACT. In this paper, we give a new generalization of positive sectional curvature called *positive weighed sectional curvature*, which depends on a choice of Riemannian metric and a smooth vector field. We give several simple examples of Riemannian metrics that do not have positive sectional curvature but support a vector field that gives them positive weighted curvature. On the other hand, we generalize a number of the foundational results for compact manifolds with positive sectional curvature to positive weighted curvature. In particular, we prove generalizations of Weinstein’s theorem, O’Neill’s formula for submersions, Frankel’s theorem, and Wilking’s connectedness lemma. As applications of these results, we recover weighted versions of topological classification results of Grove-Searle and Wilking for manifolds of high symmetry rank and positive curvature.

## 1. INTRODUCTION

Understanding Riemannian manifolds with positive sectional curvature is a deep and notoriously difficult problem in Riemannian geometry. A common approach to such problems in mathematics is to generalize a problem to a more flexible one, and then study this generalization with the hope that it will shed light on the harder, original problem. Indeed, there are a number of generalizations of positive sectional curvature that have been studied. The most obvious is non-negative sectional curvature, but other conditions such as quasi-positive or almost positive curvature have been studied in the literature (see [Zil07, KT14] and references therein).

In this paper, we propose a different approach to generalizing positive curvature, one that depends on choosing a positive, smooth density function, denoted by  $e^{-f}$ , or a smooth vector field  $X$ . Our motivation for considering such a generalization is the corresponding theory of Ricci curvature for manifolds with density,

which was studied by Lichnerowicz [Lic70, Lic72] and was later generalized and popularized by Bakry-Emery and their collaborators [BÉ85]. There are too many recent results in this area to reference all of them here, but some that are more relevant to this article include [Lot03, Mor05, Mor09b, MW12, WW09]. Also see Chapter 18 of [Mor09a] and the references therein.

For a triple  $(M^n, g, X)$ , where  $(M, g)$  is a Riemannian manifold and  $X$  is a smooth vector field, the  $m$ -Bakry-Emery Ricci tensor is

$$\text{Ric}_X^m = \text{Ric} + \frac{1}{2}L_X g - \frac{X^\# \otimes X^\#}{m},$$

where  $m$  is a constant that is also allowed to be infinite, in which case we write  $\text{Ric}_X^\infty = \text{Ric}_X = \text{Ric} + \frac{1}{2}L_X g$ . For a manifold with density, we set  $X = \nabla f$  and write  $\text{Ric}_f^m = \text{Ric} + \text{Hess } f - \text{d}f \otimes \text{d}f/m$ .

The Bakry-Emery Ricci tensors come up in many areas of geometry and analysis, including optimal transport [LV09, Stu06a, Stu06b, vRS05], the isoperimetric inequality [Mor05], and the Ricci flow [Per]. Our definition of positive weighted sectional curvature, which looks similar to the Bakry-Emery Ricci tensors, is the following.

**Definition.** A Riemannian manifold  $(M, g)$  equipped with a vector field  $X$  has *positive weighted sectional curvature* if, for every point  $p \in M$ , every 2-plane  $\sigma \in T_p M$ , and every unit vector  $V \in \sigma$ , one of the following hold:

- $\sec(\sigma) + \frac{1}{2}(L_X g)(V, V) > 0$ ;
- $X = \nabla f$  and  $\sec(\sigma) + \text{Hess } f(V, V) + \text{d}f(V)^2 > 0$  for some function  $f$ .

Note that a Riemannian manifold with positive sectional curvature admits positive weighted sectional curvature, where  $X$  is chosen to be zero. The converse to this statement does not hold, as we show by example in Propositions 3.11 and 3.16. (For additional examples that further illustrate the difference between these notions, see Section 3.)

This definition is motivated by earlier work of the second author [Wy115], where generalizations of classical results such as the classification of constant curvature spaces, the theorems of Cartan-Hadamard, Synge, and Bonnet-Myers, and the (homeomorphic) quarter-pinched sphere theorem are proven for manifolds with density.

There are a number of reasons why positive weighted sectional curvature is a natural generalization of positive sectional curvature. We will discuss this in more detail in Section 2. For example, we observe in Section 2 that the following low-dimensional result holds (see Theorem 2.1 and the following remarks). It follows from earlier work of the second author [Wy108, Wy115].

**Theorem A.** *Suppose  $M$  is a compact manifold of dimension two or three. If  $M$  admits a metric and a vector field with positive weighted sectional curvature, then  $M$  is diffeomorphic to a spherical space form.*

This raises the following motivating question in higher dimensions.

**Question (Motivating Question).** *If  $(M^n, g, X)$  is compact and with positive weighted curvature, does  $M$  admit a metric of positive sectional curvature?*

Theorem A shows the answer is “yes” in dimensions two and three. On the other hand, we show there are complete metrics with density on  $\mathbb{R} \times T^n$  with positive weighted sectional curvature. By a theorem of Gromoll-Meyer [GM69],  $\mathbb{R} \times T^n$  does not admit a metric of positive curvature, so the answer is “no” in the complete case.

We approach this question by considering spaces with a high amount of symmetry. Since the 1990s, when Grove popularized the approach, quite a lot of powerful machinery has been developed for studying manifolds with positive curvature through symmetry. (See the survey articles [Wil07, Gro09, Zil14] for details as well as many applications.)

A first consideration is that a given vector field  $X$  may not be invariant under the isometries of  $g$ . In Section 4, we deal with this issue by showing that, given a triple  $(M, g, X)$  with positive weighted curvature and a compact group of isometries  $G$  acting on  $(M, g)$ , it is always possible to change  $X$  to  $\bar{X}$  that is invariant under  $G$  so that  $(M, g, \bar{X})$  has positive weighted sectional curvature. The fact that we can always assume that the density is invariant under a fixed compact subgroup of isometries will be a key observation in most of our results. In fact, it immediately gives the following result in the homogeneous case (see Proposition 4.7).

**Theorem B.** *If a compact, homogeneous Riemannian manifold  $(M, g)$  supports a gradient field  $X = \nabla f$  such that  $(M, g, X)$  has positive weighted curvature, then  $(M, g)$  has positive sectional curvature.*

Simple examples show that this proposition is not true if the manifold is not compact (see Example 3.2). In Section 3, we also give examples of cohomogeneity-one metrics on spheres and projective spaces that have positive weighted sectional curvature but not positive sectional curvature, so the homogeneous assumption cannot be weakened.

Another way to quantify that a Riemannian manifold has a large amount of symmetry is the symmetry rank, which is the largest dimension of a torus that acts effectively on  $M$  by isometries. Our main result regarding symmetry rank and positive weighted sectional curvature is an extension of the maximal symmetry rank theorem of Grove-Searle [GS94] to positive weighted sectional curvature (see Theorem 8.1).

**Theorem C (Maximal symmetry rank theorem).** *Let  $(M^n, g, X)$  be closed with positive weighted sectional curvature. If  $T^r$  is a torus acting effectively by isometries on  $M$ , then  $r \leq \lfloor (n + 1)/2 \rfloor$ . Moreover, if equality holds and  $M$  is simply connected, then  $M$  is homeomorphic to  $\mathbb{S}^n$  or  $\mathbb{C}\mathbb{P}^{n/2}$ .*

In higher dimensions, Wilking has shown that one can assume less symmetry and still obtain a homotopy classification [Wil03, Theorem 2]. We also give an extension of this result (see Theorem 8.3).

**Theorem D (Half-maximal symmetry rank theorem).** *Let  $(M^n, g, X)$  be closed and simply connected with positive weighted sectional curvature. If  $M$  admits an effective, isometric torus action of rank  $r \geq n/4 + \log_2 n$ , then  $M$  is homeomorphic to  $S^n$  or tangentially homotopic equivalent to  $\mathbb{C}P^{n/2}$ .*

Theorems C and D show that the answer to our motivating question is “yes” (at least up to homeomorphism or homotopy) in the case of high-enough symmetry rank. On the other hand, our results are slightly weaker than the results in the unweighted setting. (We discuss this further in Sections 8 and 9.)

There are two key tools used in the proofs of Theorems C and D. The first is an extension of Berger’s theorem (Corollary 6.3) to the weighted case. The proof follows as in [GS94], and makes use of the O’Neill formula in the weighted case (Theorem 5.1). The second main tool is a generalization of Wilking’s connectedness lemma [Wil03, Theorem 2.1] to positive weighted sectional curvature (see Theorem 7.4).

**Theorem E (Wilking’s connectedness lemma).** *Let  $(M^n, g, X)$  be closed with positive weighted sectional curvature. The following hold:*

- (1) *If  $X$  is tangent to  $N^{n-k}$ , a closed, totally geodesic, embedded submanifold of  $M$ , then the inclusion  $N \rightarrow M$  is  $(n - 2k + 1)$ -connected.*
- (2) *If  $X$  and  $N^{n-k}$  are as above, and if  $G$  acts isometrically on  $M$ , fixes  $N$  pointwise, and has principal orbits of dimension  $\delta$ , then the inclusion  $N \rightarrow M$  is  $(n - 2k + 1 + \delta)$ -connected.*
- (3) *If  $X$  is tangent to  $N_1^{n-k_1}$  and  $N_2^{n-k_2}$ , a pair of closed, totally geodesic, embedded submanifolds with  $k_1 \leq k_2$ , then  $N_1 \cap N_2 \rightarrow N_2$  is  $(n - k_1 - k_2)$ -connected.*

The only assumption in Theorem E not needed in the unweighted version is that  $X$  be tangent to the submanifolds. This of course is true in the unweighted setting where  $X = 0$ . In the applications, this extra assumption holds, since the submanifolds to which we are applying the result will be fixed-point sets of isometries, and  $X$  will be invariant under these actions (see Corollary 7.2 and the following discussion). The proof of Theorem E follows from Wilking’s arguments in [Wil03] by using the second variation formula for the weighted curvatures derived in [Wy115] in place of the classical one.

This paper is organized as follows. In Sections 2 and 3, we recall the notion of weighted sectional curvature from [Wy115], define positive weighted sectional curvature, survey its basic properties (including Theorem A), and construct a number of examples. In Sections 4–7, we establish these properties and use them to prove Theorem B as well as generalizations of the O’Neill formulas, Weinstein’s theorem, and Wilking’s connectedness lemma (Theorem E). In Section 8, we use these tools to prove Theorems C and D. Finally, in Section 9, we discuss future directions.

2. DEFINITIONS AND MOTIVATION

In this section, we fix some notation and go into more detail about the motivation for the definition of positive weighted sectional curvature. At the end of this section (see Subsection 2.3), we address the fact that weighted sectional curvature is not simply a function of 2-planes in the way that sectional curvature is, and we discuss a symmetrized version of weighted sectional curvature that is such a function.

**2.1. Definition of positive weighted sectional curvature.** First, we recall some notation from [Wy115]. For a Riemannian manifold  $(M, g)$  and a vector  $V$  on  $M$ , we will call the symmetric  $(1, 1)$ -tensor  $R^V$ , given by

$$R^V(U) = R(U, V)V = \nabla_U \nabla_V V - \nabla_V \nabla_U V - \nabla_{[U, V]}V,$$

the *directional curvature operator in the direction of  $V$* . Given a smooth vector field  $X$ , the *weighted directional curvature operator in the direction of  $V$*  is another symmetric  $(1, 1)$ -tensor,

$$R_X^V = R^V + \frac{1}{2}(L_X g)(V, V) \text{ id},$$

where  $\text{id}$  is the identity operator. The *strongly weighted directional curvature operator in the direction of  $V$*  is defined as  $\bar{R}_X^V = R_X^V + g(X, V)^2 \text{id}$ .

Given an orthonormal pair  $(U, V)$  of vectors in  $T_p M$  for some  $p \in M$ , the sectional curvature  $\text{sec}(U, V)$  of the plane spanned by  $U$  and  $V$  is, by definition,  $\text{sec}(U, V) = g(R^V(U), U)$ . In the weighted cases, we similarly define

$$\begin{aligned} \text{sec}_X^V(U) &= g(R_X^V(U), U) = \text{sec}(V, U) + \frac{1}{2}(L_X g)(V, V), \\ \overline{\text{sec}}_X^V(U) &= g(\bar{R}_X^V(U), U) = \text{sec}_X^V(U) + g(X, V)^2. \end{aligned}$$

We say that  $\text{sec}_X \geq \lambda$  if  $\text{sec}_X^V(U) \geq \lambda$  for every orthonormal pair  $(V, U)$ , or, equivalently, if all of the eigenvalues of  $R_X^V$  are at least  $\lambda$  for every unit vector  $V$ . We define the condition  $\overline{\text{sec}}_X \geq \lambda$  in the analogous way. Note that  $\overline{\text{sec}}_X^V(U) \geq \text{sec}_X^V(U)$ , so that  $\text{sec}_X \geq \lambda$  implies  $\overline{\text{sec}}_X \geq \lambda$ .

In terms of this notation, we can then rephrase the definition of positive weighted sectional curvature.

**Definition.** A Riemannian manifold  $(M, g)$  equipped with a vector field  $X$  has *positive weighted sectional curvature* if one of the following hold:

- $\text{sec}_X > 0$ ;
- $X = \nabla f$  and  $\overline{\text{sec}}_f > 0$  for some function  $f$ .

Note that, unlike  $\text{sec}(U, V)$ , the weighted sectional curvatures are not symmetric in  $U$  and  $V$ . This may at first seem unnatural, but it is necessary if we want

the weighted sectional curvatures to agree with the Bakry-Emery Ricci curvatures in dimension two, as the Bakry-Emery Ricci tensors of a surface with density will generally have two different eigenvalues. (See Section 2.3 for a discussion of a symmetrized version.)

Also note that  $\sec_X^V$  and  $\overline{\sec}_X^V$  average to Bakry-Emery Ricci curvatures in the following sense. Let  $\{E_i\}_{i=1}^{n-1}$  be an orthonormal basis of the orthogonal complement of  $V$ ; then,

$$\begin{aligned}\operatorname{Ric}_{(n-1)X}(V, V) &= \sum_{i=1}^{n-1} \sec_X^V(E_i), \\ \operatorname{Ric}_{(n-1)X}^{-(n-1)}(V, V) &= \sum_{i=1}^{n-1} \overline{\sec}_X^V(E_i).\end{aligned}$$

In particular, for surfaces,  $\sec_X \geq \lambda$  is equivalent to  $\operatorname{Ric}_X \geq \lambda$ , and similarly for  $\overline{\sec}_X$  and  $\operatorname{Ric}_X^{-1}$ . The curvature  $\operatorname{Ric}_{(n-1)X}^{-(n-1)}$  is an example of Bakry-Emery Ricci curvature with negative  $m$  that has been studied recently in [KM, Oht].

**2.2. Properties of positive weighted sectional curvature.** Now that we have introduced the main equations involving weighted sectional curvature, we summarize some of the properties that the condition of positive weighted sectional curvature shares with positive sectional curvature. We then give a basic outline of how these facts lead to the proof of Theorems C and D.

First, positive weighted sectional curvature is preserved under covering maps. Specifically, if  $(M, g, X)$  has positive weighted sectional curvature and  $\tilde{M}$  is a cover of  $M$ , then  $(\tilde{M}, \tilde{g}, \tilde{X})$  has positive weighted sectional curvature where  $\tilde{g}$  and  $\tilde{X}$  are the pullbacks of  $g$  and  $X$ , respectively, under the covering map.

A second property of positive weighted sectional curvature is that the fundamental group is finite in the compact case. Indeed, this follows from [Wy108, Theorem 1.1] and [Wy115, Theorem 1.14] by the fact that positive weighted sectional curvature lifts to covers, as follows.

**Theorem 2.1.** *Let  $(M, g)$  be a complete Riemannian manifold. If there exists a vector field  $X$  such that  $\operatorname{Ric}_X > \lambda > 0$ , or if  $M$  is compact and there is a function  $f$  such that  $\operatorname{Ric}_f^{-(n-1)} > \lambda > 0$ , then  $\pi_1(M)$  is finite.*

This theorem immediately implies the classification of compact two- and three-dimensional manifolds with positive weighted sectional curvature stated in Theorem A. Indeed, this follows in dimension two from the classification of surfaces, and in dimension three from the Ricci flow proof of the Poincaré conjecture.

We note that, for positive Ricci curvature, the finiteness of fundamental group follows from the Bonnet-Myers diameter estimate. There is no diameter estimate for the weighted curvatures, as there are complete non-compact examples with  $\sec_f > \lambda > 0$  (see Example 3.2).

A third property of positive weighted sectional curvature is that the vector field  $X$  can always be chosen so that it is invariant under a fixed compact group of

isometries. We interpret this as a shared property with positive sectional curvature, since the zero vector field is always invariant. Specifically we have the following result.

**Corollary 2.2.** *If  $(M, g, X)$  has positive weighted sectional curvature, and if  $G$  is a compact subgroup of the isometry group of  $(M, g)$ , then  $X$  can be replaced by a  $G$ -invariant vector field  $\tilde{X}$  such that  $(M, g, \tilde{X})$  has positive weighted sectional curvature.*

When  $M$  is compact, the isometry group is compact, and hence this corollary applies in this case where  $G$  is the entire isometry group. As we mentioned in the [Introduction](#), reducing to the invariant case will be key in most of our results. Corollary 2.2 follows immediately from Lemmas 4.3 and 4.5 below.

A fourth property of positive weighted sectional curvature is that Riemannian submersions preserve it in the following sense.

**Corollary 2.3.** *Let  $\pi : (M, g) \rightarrow (B, h)$  be a Riemannian submersion. Let  $X$  be a vector field  $X$  on  $M$  that descends to a well-defined vector field  $\pi_*X$  on  $B$ . If  $(M, g, X)$  has positive weighted sectional curvature, then so does  $(B, h, \pi_*X)$ .*

This follows immediately from a generalization of O'Neill's formulas proved below (Theorem 5.1). We also obtain from O'Neill's formulas that Cheeger deformations preserve positive weighted sectional curvature (Lemma 5.3).

Corollary 2.3 implies the following: if  $(M, g, X)$  is compact with positive weighted sectional curvature and  $G$  is a closed subgroup of the isometry group that acts freely on  $M$ , then  $M/G$  admits positive weighted sectional curvature. Indeed, by Corollary 2.2 we can modify  $X$  so that it is  $G$ -invariant and thus descends to a vector field on  $M/G$  via the quotient map  $\pi : M \rightarrow M/G$ . It follows that  $M/G$  equipped with the vector field  $\pi_*X$  has positive weighted sectional curvature by Corollary 2.3. We implicitly use this fact in the proof of Berger's theorem (see Corollary 6.3).

Finally, a crucial property of positive weighted sectional curvature is that Synge-type arguments for positive sectional curvature generalize to the weighted setting. This follows from studying a second variation formula for energy of geodesics that was derived in [Wy115]. Given a variation  $\bar{\gamma} : [a, b] \times (-\varepsilon, \varepsilon) \rightarrow M$  of a geodesic  $\gamma = \bar{\gamma}(\cdot, 0)$ , let  $V = (\partial\bar{\gamma}/\partial s)|_{s=0}$  denote the variation vector field along  $\gamma$ . The second variation of energy is given by

$$\frac{d^2}{ds^2} \Big|_{t=0} E(\gamma_s) = I(V, V) + g \left( \frac{\partial^2 \bar{\gamma}}{\partial s^2} \right) \Big|_{t=a}^{t=b},$$

where  $I(V, V)$  is the index form of  $\gamma$ . The usual formula for the index form is

$$I(V, V) = \int_a^b (|V'|^2 - R^{y'}(V, V)) dt.$$

In terms of the weighted directional curvature operators, the index form can be re-written as follows (see [Wy115, Section 5]):

$$\begin{aligned}
 (2.1) \quad I(V, V) &= \int_a^b (|V'|^2 - R_X^{Y'}(V, V) - 2g(Y', X)g(V, V')) dt \\
 &\quad + g(Y', X)|V|^2 \Big|_{t=a}^{t=b} \\
 (2.2) \quad &= \int_a^b (|V' - g(Y', X)V|^2 - \bar{R}_X^{Y'}(V, V)) dt \\
 &\quad + g(Y', X)|V|^2 \Big|_{t=a}^{t=b}.
 \end{aligned}$$

It may not be immediately apparent why these formulas are natural, but they do allow us to generalize Synge-type arguments using the following result.

**Lemma 2.4.** *Fix a triple  $(M, g, X)$ . Let  $\gamma : [a, b] \rightarrow M$  be a geodesic on  $M$ , and let  $Y$  be a unit-length, parallel vector field along and orthogonal to  $\gamma$ . The following hold:*

- (1) *If  $\sec_X > 0$ , then the variation  $\gamma_s(t) = \exp(sY)$  of  $\gamma$  satisfies*

$$\frac{d^2}{ds^2} \Big|_{s=0} E(\gamma_s) < g(Y'(t), X_{Y(t)}) \Big|_{t=a}^{t=b}.$$

- (2) *If  $X = \nabla f$  and  $\overline{\sec}_f > 0$ , then the variation  $\gamma_s(t) = \exp(se^f Y)$  of  $\gamma$  satisfies*

$$\frac{d^2}{ds^2} \Big|_{s=0} E(\gamma_s) < e^{f(Y(t))} g(Y'(t), X_{Y(t)}) \Big|_{t=a}^{t=b}.$$

This lemma is used in Sections 6 and 7 to generalize theorems of Weinstein, Berger, Synge, and Frankel, as well as Wilking's connectedness lemma. Once we have these results, it is not hard to see how to generalize the proofs of Theorems C and D to the weighted setting. We indicate briefly how the arguments go.

The proofs proceed by induction on the dimension  $n$ , with the base cases  $n \in \{2, 3\}$  being handled by the classification of simply connected, compact manifolds in these dimensions. If a torus acts effectively on  $M$ , which has positive weighted sectional curvature, then we obtain a fixed point set  $N$  of lower dimension by Berger's theorem. The fixed point set of a subgroup of isometries is always a totally geodesic submanifold, and since we can assume  $X$  is invariant under the group, we also obtain that  $X$  is tangent to  $N$ . It follows immediately that  $N$  with restricted vector field  $X$  also has positive weighted sectional curvature. Finally, we can restrict the torus action on  $M$  to a torus action on  $N$ , so it might follow by induction on the dimension of the manifold that  $N$  satisfies the conclusion of the theorem. If the induction hypothesis does not apply, the codimension of  $N$  is small, and other arguments are used to again show that  $N$  satisfies

the conclusion of the theorem. By applying Wilking’s connectedness lemma, the topology of  $M$  is recovered from the topology of  $N$ .

**2.3. Symmetrized weighted sectional curvature.** Unlike sectional curvature, the weighted sectional curvatures do not define a function of 2-planes. In this subsection, we define a symmetrized version that does. We also compare the notions of sectional curvature and symmetrized weighted sectional curvature.

Given a vector field  $X$  on a Riemannian manifold  $M$ ,  $\text{sec}_X$  can be regarded as a function  $\text{sec}_X(\sigma, V)$  of  $(\sigma, V)$ , where  $\sigma \subseteq T_pM$  is a two-plane and  $V$  is a unit vector in  $\sigma$ . To evaluate  $\text{sec}_X(\sigma, V)$ , choose either of the two unit vectors in  $\sigma$  orthogonal to  $V$ , call it  $U$ , and evaluate  $\text{sec}_X^V(U)$ .

Note that the unit circle  $\mathbb{S}^1(\sigma)$  in  $\sigma$  is defined by the metric, so it makes sense to average over unit vectors  $e^{i\theta} \leftrightarrow V \in \mathbb{S}^1(\sigma)$ . We denote this by

$$\text{sym sec}_X(\sigma) = \frac{1}{2\pi} \int_0^{2\pi} \text{sec}_X(\sigma, e^{i\theta}) \, d\theta.$$

One can similarly define  $\text{sym } \overline{\text{sec}}_X$ . One appealing aspect of this curvature quantity is that it is the same kind of object as  $\text{sec}$ , a function on two-planes.

This definition was motivated by a suggestion of Guofang Wei, who suggested looking at the quantity

$$\text{sec}_X^V(U) + \text{sec}_X^U(V).$$

Note that  $\frac{1}{2}(\text{sec}_X^V(U) + \text{sec}_X^U(V))$  equals  $\text{sym sec}_X$ , and likewise in the strongly weighted case.

We analyze the conditions  $\text{sym sec}_X > 0$  and  $\text{sym } \overline{\text{sec}}_X > 0$  in dimension two. First, it is clear that in any dimension,

$$\begin{array}{ccc} \text{sec}_X > 0 & \Rightarrow & \text{sym sec}_X > 0 \\ \Downarrow & & \Downarrow \\ \overline{\text{sec}}_X > 0 & \Rightarrow & \text{sym } \overline{\text{sec}}_X > 0. \end{array}$$

Second, in dimension two,  $\text{sym sec}_f = \text{scal} / 2 + \Delta f$ . This is the same as the weighted Gauss curvature studied in [CHH<sup>+</sup>06, CM11], which contain proofs that the Gauss-Bonnet theorems hold for this weighted curvature. In particular, we have the following result (compare [CHH<sup>+</sup>06, Proposition 5.3]).

**Theorem 2.5 (Gauss-Bonnet).** *If  $M^2$  is orientable, then*

$$\int_M \text{sym sec}_X = 2\pi\chi(M).$$

This gives the following generalization of one case of Theorem A, which implies that a two-dimensional, compact manifold  $M$  that admits  $\text{sec}_X > 0$  for some vector field  $X$  is diffeomorphic to a spherical space form.

**Corollary 2.6.** *If  $M^2$  is compact and admits a metric and vector field  $X$  with  $\text{sym sec}_X > 0$ , then  $M^2$  is diffeomorphic to a spherical space form.*

On the other hand, the torus  $T^2$ , while it does not admit  $\text{sym sec}_X > 0$ , does admit a metric with  $\text{sym } \overline{\text{sec}}_X > 0$ . To see this, equip the torus with a flat metric and a unit-length Killing field  $X$ ; then, we have

$$\text{sym } \overline{\text{sec}}_X = 0 + 0 + \frac{1}{2\pi} \int_0^{2\pi} g(X, e^{i\theta})^2 d\theta = 1.$$

In fact, this example can be immediately generalized as follows.

**Proposition 2.7.** *If  $(N, g)$  is a Riemannian manifold with positive sectional curvature, then  $S^1 \times N$  admits a metric and a vector field  $X$  such that  $\text{sym } \overline{\text{sec}}_X > 0$ .*

*Proof.* Let  $g$  be the product metric, and let  $X$  denote the unit-length Killing field tangent to the circle factor. If  $\sigma$  is a two-plane tangent to  $N$ , then

$$\text{sym } \overline{\text{sec}}_X(\sigma) \geq \text{sec}^{g_N}(\sigma) > 0.$$

If  $\sigma$  is a two-plane not contained in the tangent space to  $N$ , then

$$\text{sym } \overline{\text{sec}}_X(\sigma) \geq \frac{1}{2} |\text{proj}_\sigma(X)|^2 > 0,$$

where  $\text{proj}_\sigma$  denotes the projection onto  $\sigma$ . □

This raises the following question.

**Question 2.8.** *Does the torus admit  $\overline{\text{sec}}_X > 0$  or  $\text{sym } \overline{\text{sec}}_f > 0$ ? More generally, are there compact manifolds with  $\overline{\text{sec}}_X > 0$  or  $\text{sym } \overline{\text{sec}}_f > 0$  and infinite fundamental group?*

We point out that Gauss-Bonnet type arguments do not seem to give a different proof that any compact surface with density with  $\overline{\text{sec}}_f > 0$  is a sphere. Indeed, if we trace  $\overline{\text{sec}}_X$ , we obtain  $\text{scal} + \text{div}(X) + |X|^2$ . The integral of this is  $4\pi\chi(M) + \int_M |X|^2 d\text{vol}_g$ , which is not a topological quantity.

We also note that the Gauss-Bonnet theorem gives interesting information about other inequalities involving curvature. First, we consider a positive lower bound.

**Proposition 2.9.** *Let  $(M, g)$  be a compact surface with  $\text{sym sec}_X \geq 1$ . The area of  $M$  is at most  $4\pi$ . Moreover, if  $\text{sec}_X \geq 1$  and the area of  $M$  is  $4\pi$ , then  $(M, g)$  is the round sphere, and  $X$  is a Killing field.*

*Proof.* We apply the discussion above to the universal cover  $\tilde{M}$  of  $M$ , endowed with the pulled back metric  $\tilde{g}$  and vector field  $\tilde{X}$ . It follows that  $\chi(\tilde{M}) > 0$ , so that  $\chi(\tilde{M}) = 2$  and  $\text{area}(M) \leq \text{area}(\tilde{M}) \leq 4\pi$ . Moreover, if  $\text{area}(M) = 4\pi$ ,

then both of these inequalities are equalities. In particular,  $\pi_1(M)$  is trivial, and  $\sec_{\tilde{X}} = \text{Ric}_{\tilde{X}} = \tilde{g}$ , so that  $(M, g, X) = (\tilde{M}, \tilde{g}, \tilde{X})$  is a compact, two-dimensional Ricci soliton. A result of Chen, Lu, and Tian [CLT06] then shows that  $M$  has constant curvature 1 and that  $X$  is a Killing field. Since  $M$  is simply connected, this proves the proposition.  $\square$

We can also consider the case of negative curvature in dimension two. It was shown in [Wy115] that if a compact manifold has  $\overline{\sec}_X \leq 0$ , then the universal cover is diffeomorphic to Euclidean space, showing that a compact surface admits  $\overline{\sec}_X \leq 0$  if and only if it is not the sphere or real projective space. In fact, the Gauss-Bonnet argument improves this result for surfaces, as it shows that the conclusion holds if  $\text{sym sec}_X \leq 0$ . Moreover, it also shows that if a metric on the torus supports a vector field with  $\text{sym sec}_X \leq 0$ , then the metric is flat and  $X$  is Killing. In particular, the two-torus has no metric with density on it with  $\text{sym sec}_X < 0$ .

The discussion above, along with the work of Corwin and Morgan [CM11] certainly shows that the study of the symmetrized weighted sectional curvature is warranted. In fact, the results in Section 5 of this paper about Riemannian submersions and Cheeger deformation have analogues for the symmetrized curvatures with the same proofs. On the other hand, there does not seem to be a good second variation formula for the symmetrized curvatures that can give us a version of Lemma 2.4. Note that the unsymmetrized curvatures also appear in the second variation of the weighted distance (see [Mor06, Mor09b]). Without some kind of second variation formula for the symmetrized curvatures, it seems unlikely that the other results of this paper can be generalized to the symmetrized case, or that many of the facts for surfaces mentioned above can be generalized to higher dimensions.

### 3. EXAMPLES

In this section, we discuss a number of examples of metrics with positive weighted curvature, including some that do not have positive sectional curvature. As a warm-up, we first consider the case of products.

**Definition 3.1.** Given  $(M_1, g_1, X_1)$  and  $(M_2, g_2, X_2)$  where  $(M_i, g_i)$  are Riemannian manifolds and  $X_i$  are smooth vector fields, the product of  $(M_1, g_1, X_1)$  and  $(M_2, g_2, X_2)$  is the triple  $(M_1 \times M_2, g_1 + g_2, X_1 + X_2)$ .

A basic fact about positive sectional curvature is that it is not preserved by taking products, as the sectional curvature of a plane spanned by vectors in each factor is zero. Indeed, one of the most famous open problems in Riemannian geometry is the Hopf conjecture, which states that  $\mathbb{S}^2 \times \mathbb{S}^2$  does not admit any metric of positive sectional curvature. In the weighted case, there are noncompact examples of products that have positive weighted sectional curvature.

**Example 3.2.** We define the one-dimensional Gaussian as the real line  $\mathbb{R}$  with coordinate  $x$ , standard metric  $g = dx^2$ , and vector field  $X = \frac{1}{2}\nabla(x^2) = x\text{d}/\text{d}x$ .

This triple has  $\text{sec}_X = 1$ . If we take the product of two one-dimensional Gaussians, we get a two-dimensional Gaussian: namely,  $\mathbb{R}^2$  with the Euclidean metric and vector field  $X = \nabla f$  where  $f(A) = \frac{1}{2}|A|^2$ . This triple still has  $\text{sec}_X = 1$ . Moreover, taking further products, we obtain the  $n$ -dimensional Gaussian as the product of  $n$  one-dimensional Gaussians, all of which have  $\text{sec}_X = 1$ .

On the other hand, it is easy to see that such examples cannot exist in the compact case.

**Proposition 3.3.** *No product of the form  $(M_1 \times M_2, g_1 + g_2, X_1 + X_2)$  with one of the  $M_i$  compact has positive weighted sectional curvature.*

*Proof.* Let  $M_1$  be the compact factor, and suppose first that  $\text{sec}_X > 0$ . Consider the “vertizontal” curvature given by  $Y$  tangent to  $M_1$  and  $U$  tangent to  $M_2$ :

$$\text{sec}_X(Y, U) = \text{sec}(Y, U) + \frac{1}{2}L_X g(Y, Y) = \frac{1}{2}L_{X_1} g_1(Y, Y).$$

This shows that if  $\text{sec}_X > 0$ , then  $\frac{1}{2}L_{X_1} g_1 > 0$ . This is impossible if  $M_1$  is compact by the divergence theorem, as  $\text{tr}(L_{X_1} g) = \text{div}(X_1)$ .

The case where  $\overline{\text{sec}}_f > 0$  is analogous. Here, we obtain that the function  $u_1 = e^{f_1}$  has  $\text{Hess}_{g_1} u_1 > 0$  on  $M_1$ , which is again impossible on a compact manifold. □

This shows that the Hopf conjecture is also interesting for weighted sectional curvature.

**Question 3.4 (Weighted Hopf conjecture).** *Does  $\mathbb{S}^2 \times \mathbb{S}^2$  admit a metric and vector field with positive weighted sectional curvature?*

In the next few sections, we investigate examples with positive weighted sectional curvature, using the simple construction of warped products over a one-dimensional base. As we can see even in the case of products, it is easier to construct non-compact examples than compact ones, so we will investigate the non-compact case first.

**3.1. Noncompact examples.** A warped product metric over a one-dimensional base is a metric of the form  $g = dr^2 + \varphi^2(r)g_N$  where  $N$  is an  $(n - 1)$ -dimensional manifold. Up to rescaling  $\varphi$  and the fiber metric  $g_N$  and reparametrizing  $r$ , there are three possibilities for the topology of complete metrics of this form:

- (1) If  $\varphi(r) > 0$  for  $r \in \mathbb{R}$  and  $(N, g_N)$  is complete, then  $g$  gives a complete metric on  $\mathbb{R} \times N$ . If  $\varphi$  is also periodic, then we can take the quotient to get a metric on  $\mathbb{S}^1 \times N$ .
- (2) If  $\varphi(r) > 0$  for  $r \in (0, \infty)$  and  $\varphi$  is an odd function with  $\varphi'(0) = 1$ , and if  $(N, g_N)$  is a round sphere of constant curvature 1, then  $g$  defines a complete rotationally symmetric metric on  $\mathbb{R}^n$ .

- (3) If  $\varphi(r) > 0$  for  $r \in (0, R)$  and  $\varphi$  is an odd function at 0 and  $R$  with  $\varphi'(0) = 1$  and  $\varphi'(R) = -1$ , and if  $(N, g_N)$  is a round sphere of constant curvature 1, then  $g$  defines a complete rotationally symmetric metric on  $\mathbb{S}^n$ .

For  $Y, Z$  tangent to  $g_N$ , we have the following well-known formulas for the curvature operator of a one-dimensional warped product:

$$\begin{aligned} \mathcal{R}(\partial_r \wedge Y) &= -\frac{\varphi''}{\varphi} \partial_r \wedge Y, \\ \mathcal{R}(Y \wedge Z) &= \mathcal{R}^N(Y \wedge Z) \left(\frac{\varphi'}{\varphi}\right)^2 Y \wedge Z, \end{aligned}$$

where  $\mathcal{R}^N$  denotes the curvature operator of  $N$ . We will be interested in lower bounds on weighted curvature of the warped product. All of our examples will also have the property that  $X = \nabla f$ , so we focus only on this case. The following lemma simplifies the problem of proving such lower bounds for warped product metrics over a one-dimensional base.

**Lemma 3.5.** *Let  $dr^2 + \varphi^2(r)g_N$  be a warped product metric, and assume  $f$  is a smooth function that only depends on  $r$ . The weighted curvature  $\text{sec}_f \geq \lambda$  if and only if*

$$\begin{aligned} \lambda &\leq \text{sec}_f^{\partial_r}(Y) = -\frac{\varphi''}{\varphi} + f'', \\ \lambda &\leq \text{sec}_f^Y(\partial_r) = -\frac{\varphi''}{\varphi} + f' \frac{\varphi'}{\varphi}, \\ \lambda &\leq \text{sec}_f^Y(Z) = \frac{\text{sec}^{g_N}(Y, Z) - (\varphi')^2}{\varphi^2} + f' \frac{\varphi'}{\varphi}, \end{aligned}$$

for all orthonormal pairs  $(Y, Z)$ , where  $Y$  and  $Z$  are tangent to  $N$ .

Similarly,  $\overline{\text{sec}}_f \geq \lambda$  if and only if these three inequalities hold with  $f'$  replaced by  $u' / u$  and  $f''$  replaced by  $u'' / u$ , where  $u = e^f$ .

This lemma implies that one can show  $\text{sec}_f \geq \lambda$  for these metrics by plugging in “test pairs” of the form  $(\partial_r, Y)$ ,  $(Y, \partial_r)$ , and  $(Y, Z)$ , where  $Y$  and  $Z$  are tangent to  $N$ . In particular, if  $\text{sec}^{g_N}$  is bounded from below, then proving  $\text{sec}_f \geq \lambda$  reduces to showing three inequalities involving the functions  $\varphi$  and  $f$ .

*Proof.* As the proof is similar, we omit the proof in the strongly weighted case. Let  $U = a \partial_r + Y$  and  $V = b \partial_r + Z$  be an arbitrary orthonormal pair of vectors, where  $Y$  and  $Z$  are tangent to  $N$ . By orthonormality,

$$|Y \wedge Z|^2 = 1 - a^2 - b^2,$$

so  $a^2 + b^2 \leq 1$ . Since  $\partial_r \wedge (aZ - bY)$  and  $Y \wedge Z$  are eigenvalues of the curvature operator, we have

$$\begin{aligned} \sec(U, V) &= \langle \mathcal{R}(U \wedge V), U \wedge V \rangle \\ &= -\frac{\varphi''}{\varphi} |\partial_r \wedge (aZ - bY)|^2 + \left( \frac{\sec^N(Y, Z) - (\varphi')^2}{\varphi^2} \right) |Y \wedge Z|^2 \\ &= -\frac{\varphi''}{\varphi} (a^2 + b^2) + \left( \frac{\sec^N(Y, Z) - (\varphi')^2}{\varphi^2} \right) (1 - a^2 - b^2). \end{aligned}$$

Next, we calculate

$$\text{Hess } f(U, U)|V|^2 = a^2 f'' + (1 - a^2) f' \frac{\varphi'}{\varphi}.$$

Observe that  $\sec_f^U(V) = \sec(U, V) + \text{Hess } f(U, U)|V|^2$  is a linear function in the quantities  $a^2$  and  $a^2 + b^2$ . Moreover, these quantities vary over a triangle, since

$$0 \leq a^2 \leq a^2 + b^2 \leq 1,$$

so the minimal (and maximal) values of  $\sec_f^U(V)$  occur at one of the three corners. This proves the lemma, since these corners correspond to orthonormal pairs of the form  $(\partial_r, Y)$ ,  $(Y, \partial_r)$ , and  $(Y, Z)$ . □

As a first application of Lemma 3.5, we consider the problem of prescribing positive weighted sectional curvature locally on a subset of the round sphere.

**Proposition 3.6.** *Let  $M$  be a round sphere of constant curvature 1, and  $H^+$  an open round hemisphere in  $M$ . For any  $\lambda \in \mathbb{R}$ , there is a density on  $H^+$  with  $\sec_f \geq \lambda$ , and there is no density defined on an open set containing the closure of  $H^+$  with  $\overline{\sec}_f > 1$ .*

*Proof.* First, we prove the non-existence. It suffices to show that a geodesic ball  $B$  of radius  $\pi/2 + \varepsilon$  cannot admit a density  $f$  such that  $\overline{\sec}_f > 1$ . On  $B$ , we can write the round metric as the warped product  $dr^2 + \sin^2(r)g_{\mathbb{S}^{n-1}}$ , where  $r \in (0, \pi/2 + \varepsilon)$ . By Lemma 4.5 proved in the next section, we can assume that  $f = f(r)$ . By Lemma 3.5,  $\overline{\sec}_f > 1$  only if  $(u'/u) \cot(r) > 0$ . However,  $\cot(\pi/2) = 0$ , so the second inequality is impossible to satisfy.

On the other hand, to find a density  $f$  with  $\sec_f \geq \lambda$ , we only need that

$$f'' \geq \lambda - 1 \quad \text{and} \quad f' \cot(r) \geq \lambda - 1.$$

Such a density exists; for example,  $f$  given by

$$f(r) = (\lambda - 1) \int \tan(x) dx = -(\lambda - 1) \log(\cos(r))$$

satisfies these properties. Note that in these examples,  $f$  blows up at the equator  $r = \pi/2$ . □

On the other hand, we note the general fact that every point  $p$  in a Riemannian manifold has a neighborhood  $U$  supporting a density such that  $\text{sec}_f \geq \lambda$ .

**Proposition 3.7.** *Let  $(M, g)$  be a Riemannian manifold,  $p \in M$ , and  $\lambda \in \mathbb{R}$ . There exists an open set  $U$  containing  $p$  that supports a density  $f$  such that  $\text{sec}_f \geq \lambda$  on  $U$ .*

*Proof.* Let  $r$  be the distance function to  $p$ . Since  $\text{Hess } r \sim 1/r$  as  $r \rightarrow 0$ , there exists  $0 < \varepsilon < 1$  such that  $r \text{ Hess } r > \varepsilon g$  on  $B(p, \varepsilon)$ . Let  $\rho = \inf \text{sec}(B(p, \varepsilon), g)$ . Define  $f = ((\lambda - \rho)/(2\varepsilon))r^2$ . We have that

$$\text{Hess } f = \frac{\lambda - \rho}{\varepsilon} dr \otimes dr + \frac{\lambda - \rho}{\varepsilon} r \text{ Hess } r \geq (\lambda - \rho)g,$$

which implies that  $\text{sec}_f \geq \lambda$  on  $B(p, \varepsilon)$ . □

We now come to our first complete example.

**Proposition 3.8.** *Let  $(N, g_N)$  be a metric of non-negative sectional curvature. For any  $\lambda$ , the metric  $g = dr^2 + e^{2r} g_N$  on  $\mathbb{R} \times N$  admits a density of the form  $f = f(r)$  such that  $\overline{\text{sec}}_f \geq \lambda$ . On the other hand,  $g$  admits no density of the form  $f = f(r)$  with  $\text{sec}_f \geq -1 + \varepsilon$  with  $\varepsilon > 0$ .*

*Proof.* First set  $\varphi(r) = e^r$ . Because  $N$  has non-negative sectional curvature, Lemma 3.5 implies  $\overline{\text{sec}}_f \geq \lambda$  if and only if  $-1 + u''/u \geq \lambda$  and  $-1 + u'/u \geq \lambda$ . This can be achieved by taking  $u = e^{Ar}$  for some sufficiently large  $A \in \mathbb{R}$ .

On the other hand, for a general  $f = f(r)$ , if we have  $\text{sec}_f \geq -1 + \varepsilon$ , then  $f$  satisfies  $f''(r) \geq \varepsilon$  and  $f'(r) \geq \varepsilon$  for all  $r \in \mathbb{R}$ . This is impossible. □

**Remark 3.9.** Gromoll and Meyer [GM69] proved that a non-compact, complete manifold with  $\text{sec} > 0$  is diffeomorphic to Euclidean space. These examples show this is not true for  $\overline{\text{sec}}_f > 0$ . Moreover, Cheeger and Gromoll [CG72] showed that a non-compact complete manifold with  $\text{sec} \geq 0$  is the normal bundle over a compact totally geodesic submanifold called a *soul*. While our examples are topologically  $\mathbb{R} \times N$ , we note that the cross sections  $\{r_0\} \times N$  are not geometrically a “soul,” as they are not totally geodesic.

**Remark 3.10.** If we take  $g_N$  to be a flat metric, then the metric given by  $g = dr^2 + e^{2r} g_N$  is a hyperbolic metric. If we also choose  $f(r) = r$ , then we get a density with constant  $\overline{\text{sec}}_f = 0$ .

**3.2. Compact examples.** We now give examples of rotationally symmetric metrics on the  $n$ -sphere that admit a density  $f$  such that  $\text{sec}_f > 0$ , but that do not have  $\text{sec} \geq 0$ .

In general, a rotationally symmetric on the sphere will be of the form  $g = dr^2 + \varphi^2(r)g_{\mathbb{S}^{n-1}}$  for  $r \in [0, 2L]$ . The smoothness conditions for the warping function  $\varphi$  and density function  $f$  are that  $\varphi(0) = \varphi(2L) = 0$ ,  $\varphi'(0) = 1$ ,  $\varphi'(2L) = -1$ ,  $\varphi^{(\text{even})}(0) = \varphi^{(\text{even})}(2L) = 0$ , and  $f'(0) = f'(2L) = 0$ . Our main construction is contained in the following proposition.

**Proposition 3.11.** *There are rotationally symmetric metrics on  $S^n$  that support a density  $f$  such that  $\text{sec}_f > 0$ , but that do not have  $\text{sec} \geq 0$ .*

*Proof.* First, we shall define  $\varphi(r) = r$  on  $[0, \pi/6]$  and  $\varphi(r) = \sin(r)$  on  $[\pi/3, \pi/2]$ . On the interval  $(\pi/6, \pi/3)$ , extend  $\varphi$  smoothly so that  $\varphi'' \leq 0$  and  $\varphi' \geq 0$ . Then, we reflect  $\varphi$  across  $\pi/2$  to obtain a warping function defined on  $[0, \pi]$  that gives a smooth rotationally symmetric metric on the sphere. Geometrically, this metric consists of two flat discs connected by a region of positive curvature.

Now, define  $f(r) = \frac{1}{2}r^2$  on  $[0, \pi/3]$ , and extend  $f$  on  $(\pi/3, \pi/2]$  so that  $f' > 0$  on  $(\pi/3, \pi/2)$  and has  $f^{(\text{odd})}(\pi/2) = 0$ , so that  $f$  also defines a smooth function when reflected across  $\pi/2$ .

We now consider the potential function  $\lambda f''$  for a positive constant  $\lambda$ . The table below shows the values for the eigenvalues of the curvature operator and Hessian of  $\lambda f$  on the different regions:

	$-\varphi''/\varphi$	$(1 - (\varphi')^2)/\varphi^2$	$\lambda f''$	$\lambda f' \varphi'/\varphi$
$[0, \pi/6]$	0	0	$\lambda$	$\lambda$
$(\pi/6, \pi/3]$	$> 0$	$> 0$	$\lambda$	$\geq \lambda$
$(\pi/3, \pi/2]$	1	1	$\lambda f''$	$\geq 0$

By Lemma 3.5,  $\text{sec}_f \geq \lambda$  on  $[0, \pi/3]$ . On  $(\pi/3, \pi/2]$ , note that  $f'' < 0$  somewhere since  $f'$  must decrease from  $\pi/3$  to 0. However, by choosing  $\lambda$  small enough, we can make  $1 + \lambda f'' \geq \lambda$  on  $[\pi/3, \pi/2]$ , and then we will have  $\text{sec}_f \geq \lambda$  everywhere.

We have thus constructed examples with  $\text{sec}_f > 0$  but that do not have  $\text{sec} > 0$ . Of course, this example does have  $\text{sec} \geq 0$ . However, since having  $\text{sec}_f > 0$  is an open condition, we can perturb the metric in an arbitrary small way and still have  $\text{sec}_f > 0$ . This will give metrics with some negative sectional curvatures that still have  $\text{sec}_f > 0$ . □

On the other hand, we note that most rotationally symmetric metrics on the sphere do not have any density such that  $\overline{\text{sec}}_f > 0$ .

**Proposition 3.12.** *Let  $g = dr^2 + \varphi^2(r)g_{S^{n-1}}$ ,  $r \in [0, 2L]$ , be a metric on  $S^n$ . The following hold:*

- (1) *If there is a density  $f$  such that  $\text{sec}_f > 0$ , then*

$$\int_0^{2L} \frac{-\varphi''(r)}{\varphi(r)} dr \geq 0.$$

- (2) *If there is a density  $f$  such that  $\overline{\text{sec}}_f > 0$ , then  $\varphi$  has a unique critical point  $t_0$ . Moreover, at  $t_0$ , the metric has positive sectional curvature.*

*Proof.* By Lemmas 4.3 and 4.5, we can assume in either case that  $f$  is a function of  $r$ . Both results are simple consequences of the equations for curvature. For the first, we consider the equation

$$\sec_f^{\partial_r}(Y) = \frac{-\varphi''}{\varphi} + f'' > 0.$$

For  $f$  to define a smooth function, we must have  $f'(0) = f'(2L) = 0$ , so integrating the equation gives (1). In dimension two, this is the Gauss-Bonnet Theorem (Theorem 2.5) that we discussed in Section 2.3.

For (2), consider a point where  $\varphi'(t) = 0$ . Fix at this point an orthonormal pair of vectors  $Y$  and  $Z$  that are tangent to  $N$ . Since we have that  $\text{Hess } u(Y, Y) = u'(\varphi'/\varphi)g(Y, Y) = 0$ , the only way  $\overline{\text{sec}}_f^Y(\partial_r)$  and  $\overline{\text{sec}}_f^Y(Z)$  can be positive is if  $\sec(\partial_r, Y)$  and  $\sec(Y, Z)$  are positive. It follows that all sectional curvatures are positive at this point. Moreover, it follows that  $\varphi''(t) < 0$  at each critical point  $t$ , so there can be at most one critical point of  $\varphi$ .  $\square$

**Remark 3.13.** Proposition 3.12 (2), shows that a spherical “dumbbell” metric consisting of two spheres connected by a long neck of non-positive curvature does not have any density with  $\overline{\text{sec}}_f > 0$ .

We now consider doubly warped products of the form

$$g = dr^2 + \varphi^2(r)g_{\mathbb{S}^k} + \psi^2(r)g_{\mathbb{S}^m}, \quad r \in [0, L].$$

These metrics are also cohomogeneity-one with  $G = O(k + 1) \times O(m + 1)$ , so by Lemmas 4.3 and 4.5, we can assume that the density is of the form  $f = f(r)$ . We also have  $\text{Hess } r = \varphi' \varphi g_{\mathbb{S}^k} + \psi' \psi g_{\mathbb{S}^m}$ . Thus,

$$\text{Hess } f = f'' dr^2 + f' \varphi' \varphi g_{\mathbb{S}^k} + f' \psi' \psi g_{\mathbb{S}^m}.$$

In order for  $f$  to be  $C^2$ , we thus need  $f'(0) = f'(L) = 0$ .

We let  $Y, Z$  denote vectors in the  $\mathbb{S}^k$  factor, and  $U, V$  be vectors in the  $\mathbb{S}^m$  factor. The curvature operator in this case is

$$\begin{aligned} \mathcal{R}(\partial_r \wedge Y) &= -\frac{\varphi''}{\varphi} \partial_r \wedge Y, \\ \mathcal{R}(\partial_r \wedge U) &= -\frac{\psi''}{\psi} \partial_r \wedge U, \\ \mathcal{R}(Y \wedge Z) &= \frac{1 - (\varphi')^2}{\varphi^2} Y \wedge Z, \\ \mathcal{R}(U \wedge V) &= \frac{1 - (\psi')^2}{\psi^2} U \wedge V, \\ \mathcal{R}(Y \wedge U) &= -\frac{\varphi' \psi'}{\varphi \psi} Y \wedge U. \end{aligned}$$

This shows that, at a point  $(r, p, q)$ , there exists a basis  $\{E_i\}$  of the tangent space such that the following hold:

- The  $E_i$  are eigenvectors of  $\text{Hess } f$ .
- The  $E_i \wedge E_j$  for  $i < j$  are eigenvectors of  $\mathcal{R}$ .

In this setting, we will use the following algebraic lemma to show that certain doubly warped products on the sphere have positive weighted sectional curvature. The proof is algebraic, and is postponed until the next subsection.

**Corollary 3.14.** *Let  $(M, g)$  be a closed Riemannian manifold with non-negative curvature operator  $\mathcal{R}$ . Let  $X$  be a vector field on  $M$ . Assume that, for all  $p \in M$ , the tangent space at  $p$  has a basis  $\{E_i\}$  such that all of the following hold:*

- $E_i$  is an eigenvector for  $L_X g$  with eigenvalue  $\mu_i$  for all  $i$ .
- $E_i \wedge E_j$  is an eigenvalue for  $\mathcal{R}$  with eigenvalue  $\lambda_{ij}$  for all  $i < j$ .
- $\lambda_{ij} > 0$  or  $\min(\mu_i, \mu_j) > 0$  for all  $i < j$ .

*There exists  $\lambda > 0$  such that  $(M, g, \lambda X)$  has positive weighted sectional curvature.*

More geometrically, this result allows us to conclude that  $\text{sec}_{\lambda f} > 0$  by testing this condition on orthonormal pairs of the form  $(E_i, E_j)$  or  $(E_j, E_i)$  with  $i < j$ .

**Proposition 3.15.** *For any positive integers  $m$  and  $k$ , there is a doubly warped product metric on  $\mathbb{S}^{k+m+1}$  of the form  $g = dr^2 + \varphi^2(r)g_{\mathbb{S}^k} + \psi^2(r)g_{\mathbb{S}^m}$  with  $\text{sec}_f > 0$ , but that does not have  $\text{sec} \geq 0$ .*

*Proof.* Let  $r$  vary over the interval  $[0, \pi/2]$ , choose  $\varphi$  and  $f$  as in the proof of Proposition 3.11, and set  $\psi(r) = \cos(r)$ . The proof of Proposition 3.11 shows that we can scale  $f$  so that the weighted sectional curvatures of the pairs involving  $\partial_r$  and  $Y$  are positive. For this argument, we apply Corollary 3.14.

Choose an orthonormal basis  $\{E_i\}_{i=0}^{k+m}$  for the tangent space with  $E_0 = \partial_r$ , with  $E_1, \dots, E_k$  tangent to  $\mathbb{S}^k$ , and with  $E_{k+1}, \dots, E_{k+m}$  tangent to  $\mathbb{S}^m$ . This basis satisfies the first two conditions of Corollary 3.14. It suffices to check the third condition.

Using the expressions above for the curvature operator, all  $\lambda_{ij} > 0$ , except in the case where  $r \in [0, \pi/6]$ , and where  $E_i$  and  $E_j$  are tangent to the  $\mathbb{S}^k$  factor. For these indices, however,  $\mu_i = \text{Hess } f(E_i, E_i) > 0$  and  $\mu_j = \text{Hess } f(E_j, E_j) > 0$ . By Corollary 3.14, we have  $\text{sec}_{\lambda f} > 0$  for some  $\lambda > 0$ . The fact that we can make  $\text{sec} < 0$  for some two-planes follows for the same reason it was true in the rotationally symmetric case. □

Applying O’Neill’s formula from Section 5, this also gives us an example on  $\mathbb{C}\mathbb{P}^n$ .

**Proposition 3.16.** *There are cohomogeneity-one metrics on  $\mathbb{C}\mathbb{P}^n$  that admit a density such that  $\text{sec}_f > 0$ , but that do not have  $\text{sec} \geq 0$ .*

*Proof.* Consider a double warped product metric on the sphere  $\mathbb{S}^{2n+1}$  of the form

$$g = dr^2 + \varphi^2(r)g_{\mathbb{S}^{2n-1}} + \psi^2(r) d\theta^2.$$

Consider the Hopf fibration on  $\mathbb{S}^{2n-1}$ , and write the metric  $g_{\mathbb{S}^{2n-1}} = k + h$  where  $h$  is the metric tangent to the Hopf fiber and  $k$  is the metric on the orthogonal complement. Complex multiplication on the  $\mathbb{S}^{2n-1}$  and  $\mathbb{S}^1$  factors induces a free isometric action on  $g$ , and the quotient is  $\mathbb{C}\mathbb{P}^n$ . The quotient map is a Riemannian submersion if we equip  $\mathbb{C}\mathbb{P}^n$  with the metric

$$dr^2 + \varphi^2(r)k + \frac{(\varphi(r)\psi(r))^2}{\varphi^2(r) + \psi^2(r)}h.$$

By O'Neill's formula (Theorem 5.1), we know this metric also has  $\text{sec}_f > 0$ . Note also that if  $Y$  is a horizontal vector field in the  $\mathbb{S}^{2n-1}$  factor, then for  $r > 0$ ,  $[\partial_r, Y] = 0$ , which implies that the sectional curvature  $\text{sec}_f^{\partial_r}(Y)$  does not change under the submersion. Since there are curvatures in the doubly warped product of this form that are negative, we also obtain that the metric on  $\mathbb{C}\mathbb{P}^n$  has some negative sectional curvatures.  $\square$

**3.3. Proof of Corollary 3.14.** This section is devoted to the proof of Corollary 3.14, which is applied in the previous section. The proof is algebraic and not required for the rest of the paper, so the reader may choose to skip this subsection. The result is restated here for convenience.

**Corollary (Corollary 3.14).** *Let  $(M, g)$  be a closed Riemannian manifold with non-negative curvature operator  $\mathcal{R}$ . Let  $X$  be a vector field on  $M$ . Assume that, for all  $p \in M$ , the tangent space at  $p$  has a basis  $\{E_i\}$  such that all of the following hold:*

- $E_i$  is an eigenvector for  $L_X g$  with eigenvalue  $\mu_i$  for all  $i$ .
- $E_i \wedge E_j$  is an eigenvalue for  $\mathcal{R}$  with eigenvalue  $\lambda_{ij}$  for all  $i < j$ .
- $\lambda_{ij} > 0$  or  $\min(\mu_i, \mu_j) > 0$  for all  $i < j$ .

*There exists  $\lambda > 0$  such that  $(M, g, \lambda X)$  has positive weighted sectional curvature.*

To prove this result, first note that it suffices to prove that a  $\lambda > 0$  as in the conclusion exists at every point in  $M$ . It is then straightforward to conclude this pointwise claim from the following lemma together with the non-negativity of the curvature operator.

**Lemma 3.17.** *Let  $(V, \langle \cdot, \cdot \rangle)$  be a finite-dimensional inner product space. Let  $\mathcal{L}$  and  $\mathcal{R}$  be symmetric, linear maps on  $V$  and  $\Lambda^2 V$ , respectively. Assume there exists an orthonormal eigenbasis  $\{E_i\}$  for  $\mathcal{L}$  such that  $\{E_i \wedge E_j\}_{i < j}$  is an eigenbasis for  $\mathcal{R}$ . Denote the corresponding eigenvalues by  $\mu_i$  and  $\lambda_{ij}$ , respectively. Set  $\lambda_{ji} = \lambda_{ij}$  for  $i < j$ . Considered as a function of orthonormal pairs  $(Y, Z)$  in  $V$ , the minimum and maximum values of*

$$S(Y, Z) = \langle \mathcal{R}(Y \wedge Z), Y \wedge Z \rangle + \langle \mathcal{L}(Y), Y \rangle$$

*lie in the set*

$$\{\lambda_{ij} + \mu_i \mid i, j \text{ distinct}\} \cup \left\{ \frac{1}{2}(\lambda_{ij} + \lambda_{k\ell} + \mu_i + \mu_j) \mid i, j, k, \ell \text{ distinct} \right\}.$$

*Proof.* Let  $n = \dim(V)$ . Let  $Y = \sum a_i E_i$  and  $Z = \sum b_i E_i$  be an orthonormal pair in  $V$ . Observe that

$$S(Y, Z) = \sum_{i < j} \lambda_{ij} z_{ij} + \sum_i \mu_i x_i = S(x_i, z_{ij}),$$

where  $x_i = a_i^2$  for  $1 \leq i \leq n$ , where  $z_{ij} = (a_i b_j - a_j b_i)^2$  for  $1 \leq i < j \leq n$ . To simplify notation later, set  $z_{ii} = 0$  and  $z_{ji} = z_{ij}$  for  $1 \leq i < j \leq n$ . By orthonormality of  $(Y, Z)$ , all of the following hold:

- (1)  $x_i \geq 0$  and  $\sum x_i = 1$ , and hence the vector  $x = (x_i)$  lies on the standard simplex  $\Delta^{n-1} \subseteq \mathbb{R}^n$ .
- (2) Likewise,  $z = (z_{ij})$  lies on the standard simplex  $\Delta^{\binom{n}{2}-1} \subseteq \mathbb{R}^{n(n-1)/2}$ .
- (3) For all  $1 \leq i \leq n$ ,  $x_i \leq \sum_{j=1}^n z_{ij}$ .

Hence,  $S(Y, Z)$  equals  $S(x, z)$  for some point  $(x, z)$  in the convex polytope  $C \subseteq \mathbb{R}^n \times \mathbb{R}^{n(n-1)/2}$  defined by

$$C = \left\{ (x, z) \in \Delta^{n-1} \times \Delta^{\binom{n}{2}-1} \mid x_i \leq \sum_j z_{ij} \text{ for all } i \right\}.$$

To prove the lemma, it suffices to show that the function  $S : C \rightarrow \mathbb{R}$  has extremal values in the set described in the conclusion of the lemma.

We prove this claim by induction over  $n$ . First, if  $n = 2$ , then  $C = \Delta^1 \times \Delta^0$ , so

$$S(x, z) = \lambda_{12} + \mu_1 x_1 + \mu_2 x_2$$

has extremal values  $\lambda_{12} + \mu_1$  and  $\lambda_{12} + \mu_2$ , as claimed. Assume now that  $n \geq 3$  and that the claim holds in dimension  $n - 1$ .

Since  $C$  is a convex polytope—that is, an intersection of half-spaces—and since  $S$  is linear, the extremal values are attained at the corners (or zero-dimensional faces) of  $C$ . We now evaluate  $S$  at these corners.

Let  $(x, z) \in C$  be a corner. There exist  $0 \leq k \leq n$  and distinct indices  $i_1, \dots, i_k$  such that all of the following hold:

- (1)  $(x, z)$  lies in the interior of a  $k$ -dimensional face of  $\Delta^{n-1} \times \Delta^{\binom{n}{2}-1}$ .
- (2)  $x_{i_h} = \sum_{j=1}^n z_{i_h j}$  for  $1 \leq h \leq k$ .
- (3)  $x_i \leq \sum_{j=1}^n z_{ij}$  for all  $1 \leq i \leq n$ .

Indeed, each corner of  $C$  is obtained by intersecting some  $k$ -dimensional face of  $\Delta^{n-1} \times \Delta^{\binom{n}{2}}$  with some choice of  $k$  hyperplanes  $x_i = \sum_j z_{ij}$ . Recall that a  $k$ -dimensional face of the product is a product of an  $\ell$ -dimensional face with a  $(k - \ell)$ -dimensional face for some  $0 \leq \ell \leq k$ . Also recall that a  $k$ -dimensional face of a standard simplex is given by a choice of  $k + 1$  indices  $i_0, \dots, i_k$  for which  $x_{i_0} + \dots + x_{i_k} = 1$  and all other  $x_i = 0$ . Moreover, the interior of this face is the set of such points where, in addition, each of the  $x_{i_h} > 0$ .

First, suppose that  $k = 0$ . In other words, suppose that  $(x, z)$  lies on a corner of  $\Delta^{n-1} \times \Delta^{\binom{n}{2}}$ . There exist  $i$  and  $p < q$  such that  $x_i = 1, z_{pq} = 1$ , and all other entries of  $x$  and  $z$  are zero. By condition (3),  $i \in \{p, q\}$ , and hence  $S(x, z)$  equals  $\lambda_{iq} + \mu_i$  or  $\lambda_{pi} + \mu_i$ , as required.

Second, suppose that  $k \geq 1$  and there exists  $i_h$  with  $x_{i_h} = 0$ . By conditions (1) and (2),  $z_{i_h j} = 0$  for all  $j$ . Hence,  $S(x, z)$  does not contain any terms with index  $i_h$ . The claim follows in this case by the induction hypothesis.

Finally, suppose that  $k \geq 1$  and  $x_{i_h} > 0$  for all  $1 \leq h \leq k$ . In particular,  $x$  does not lie in a face of dimension less than  $k - 1$ . Hence, condition (1) implies that  $x$  lies in the interior of a  $(k - 1)$ - or  $k$ -dimensional face of  $\Delta^{n-1}$ , and that  $z \in \Delta^{\binom{n}{2}-1}$  lies in the interior of a one-dimensional face or a corner, respectively. We consider these cases separately:

(a) In the first case, there exists  $i_0 \notin \{i_1, \dots, i_k\}$  such that we have  $x_{i_0} > 0$  and  $x_{i_0} + x_{i_1} + \dots + x_{i_k} = 1$ . Moreover, there exists  $p < q$  such that  $z_{pq} = 1$  and  $z_{rs} = 0$  for all  $(r, s) \neq (p, q)$ . By condition (3),  $i_0 \in \{p, q\}$  and likewise for all of the distinct indices  $i_0, i_1, \dots, i_k$ . It follows that  $k$  cannot be larger than one. Moreover, if  $k = 1$ , then  $\{i_0, i_1\} = \{p, q\}$ , so

$$S(x, z) = \lambda_{i_0 i_1} + \mu_{i_0} x_{i_0} + \mu_{i_1} x_{i_1}.$$

Since  $x_{i_0}$  and  $x_{i_1}$  are positive and sum to one, this quantity is at least  $\lambda_{i_0 i_1} + \mu_{i_0}$  or  $\lambda_{i_0 i_1} + \mu_{i_1}$ , as required.

(b) In the second case,  $x_{i_1} + \dots + x_{i_k} = 1$ , and there exists  $p < q$  and  $r < s$  such that  $z_{pq} > 0, z_{rs} > 0$ , and  $z_{pq} + z_{rs} = 1$ . By condition (3),  $i_h \in \{p, q\} \cup \{r, s\}$  for all  $h$ , so clearly  $k \leq 4$ . In fact, if  $k \geq 3$ , then there exist  $i_{h_1} \in \{p, q\}$  and  $i_{h_2} \in \{r, s\}$ , which implies

$$1 = \sum_{h=1}^k x_{i_h} > x_{i_{h_1}} + x_{i_{h_2}} = \sum_j z_{i_{h_1} j} + \sum_j z_{i_{h_2} j} \geq z_{pq} + z_{rs} = 1,$$

a contradiction.

This leaves the possibilities that  $k = 2$  and  $k = 1$ . First, suppose  $k = 1$ . It follows that  $x_{i_1} = 1$  and that

$$S(x, z) = \lambda_{pq} z_{pq} + \lambda_{rs} z_{rs} + \mu_{i_1}.$$

Hence,  $S(x, z)$  is bounded between  $\lambda_{pq} + \mu_{i_1}$  and  $\lambda_{rs} + \mu_{i_1}$ . Moreover,

$$1 = x_{i_1} = \sum_j z_{i_1 j},$$

so all  $z_{ij}$  that do not appear in this sum are zero. In particular,  $i_1 \in \{p, q\}$  and  $i_1 \in \{r, s\}$ , so the claim follows in this case.

This leaves the case with  $k = 2$ . We start by showing that  $i_1$  cannot be in both  $\{p, q\}$  and  $\{r, s\}$ . Indeed, if it were, then conditions (1) and (2) imply that

$$1 = x_{i_1} + x_{i_2} > x_{i_1} = \sum_j z_{i_1 j} \geq z_{pq} + z_{rs} = 1,$$

a contradiction. By a similar argument,  $i_2$  cannot be in both sets. Condition (3) implies  $i_1, i_2 \in \{p, q\} \cup \{r, s\}$ . If  $i_1$  and  $i_2$  lie in different sets (e.g.,  $i_1 \in \{p, q\}$  and  $i_2 \in \{r, s\}$ ), then condition (2) further implies that  $x_{i_1} = z_{pq}$  and  $x_{i_2} = z_{rs}$ ; hence,

$$S(x, z) = (\lambda_{pq} + \mu_{i_1})z_{pq} + (\lambda_{rs} + \mu_{i_2})z_{rs},$$

so the claim follows in this case. Finally, if  $i_1$  and  $i_2$  lie in the same set (e.g.,  $\{p, q\}$ ), then

$$S(x, z) = \lambda_{i_1 i_2} z_{i_1 i_2} + \lambda_{rs} z_{rs} + \mu_{i_1} x_{i_1} + \mu_{i_2} x_{i_2}.$$

Moreover, in this case, condition (2) implies  $x_{i_1} = z_{pq} = x_{i_2}$ , and condition (1) implies that  $1 = x_{i_1} + x_{i_2} = 2z_{pq}$ ; hence, all four variables are equal to  $\frac{1}{2}$ . This concludes the proof of the claim.

This shows in all cases that the extremal values of  $S : C \rightarrow \mathbb{R}$  are given as in the conclusion of the lemma. As established at the beginning of the proof, the same holds true of  $S$ . □

Regarding the proof of Lemma 3.17, we note here that the point  $(x, z)$  with  $x_1 = x_2 = z_{12} = z_{34} = \frac{1}{2}$  and all other entries zero lies in the set  $C$ . Moreover, since the  $\lambda_{ij}$  and  $\mu_i$  are arbitrary, we have provided the optimal solution to the optimization problem for the function  $S : C \rightarrow \mathbb{R}$ . On the other hand,  $S(Y, Z)$  actually equals  $S(x, z)$  for some  $(x, z) \in C_0$ , where  $C_0$  is a proper subset of  $C$ . Indeed, given the definitions of  $x_i$  and  $z_{ij}$  as in the proof, it is straightforward to check that

$$(4) \quad z_{ij} \leq x_i + x_j \text{ for all } i < j.$$

Note that the point  $(x, z)$  with  $x_1 = x_2 = z_{12} = z_{34} = \frac{1}{2}$  is not in the smaller set  $C_0$ . This suggests that Lemma 3.17 could be improved to state that the optimal values are of the form  $\lambda_{ij} + \mu_i$  or  $\lambda_{ij} + \mu_j$  with  $i < j$ . Since this is not needed for our applications, we do not pursue it here.

#### 4. AVERAGING THE DENSITY

In this section, we begin to establish the properties of positive weighted sectional curvature described in Section 2. Our first consideration is that, in studying manifolds with density and symmetry, a symmetry of the metric might not be a symmetry of the density. We prove in this section that this difficulty can be overcome in the compact case. At the end, we apply these ideas to study weighted curvature properties of homogeneous metrics.

**4.1. Preservation of weighted curvature bounds under averaging.** Fix a Riemannian manifold  $(M, g)$  and a vector field  $X$  on  $M$ . Let  $G$  be a compact subgroup of the isometry group, and let  $d\mu$  denote a unit-volume, bi-invariant measure on  $G$ . Define a new,  $G$ -invariant vector field  $\bar{X}$  on  $M$  as follows:

$$\bar{X}_p = \int_G \varphi_*^{-1}(X_{\varphi(p)}) \, d\mu,$$

where we identify the elements  $\varphi \in G$  with isometries  $\varphi: M \rightarrow M$ . In the gradient case, where  $X = \nabla f$ , we similarly define  $\bar{f}(p) = \int_G f(\varphi(p)) \, d\mu$ .

As a basic observation note that, for a fixed vector field  $V$  in  $T_pM$ ,

$$\begin{aligned} g(\bar{X}, V) &= g\left(\int_G \varphi_*^{-1}(X) \, d\mu, V\right) = \int_G g(\varphi_*^{-1}(X), V) \, d\mu, \\ D_V\left(\int_G g(\varphi_*^{-1}(X), V) \, d\mu\right) &= \int_G g(\nabla_V \varphi_*^{-1}(X), V) \, d\mu \\ &\quad + \int_G g(\varphi_*^{-1}(X), \nabla_V V) \, d\mu. \end{aligned}$$

This follows from the fact that all of the functions involved are smooth, from the linearity of the integral, and from the fact that  $G$  as a compact space admits a finite partition of unity. Similar identities for passing integrals over  $G$  past a derivative also hold for the same reasons. We will use these facts repeatedly below without further comment.

We now claim the following result.

**Lemma 4.1.** *With the notation above, for any vector field  $X$  and any  $V \in T_pM$ ,*

$$(L_{\bar{X}}g)(V, V) = \int_G (L_Xg)(\varphi_*V, \varphi_*V) \, d\mu.$$

*Proof.* This follows from a straightforward calculation:

$$\begin{aligned} g(\nabla_V \bar{X}, V) &= D_V g(\bar{X}, V) - g(\bar{X}, \nabla_V V) \\ &= D_V g\left(\int_G \varphi_*^{-1}(X) \, d\mu, V\right) - g\left(\int_G \varphi_*^{-1}(X) \, d\mu, \nabla_V V\right) \\ &= \int_G D_V g(\varphi_*^{-1}(X), V) \, d\mu - \int_G g(\varphi_*^{-1}(X), \nabla_V V) \, d\mu \\ &= \int_G g(\nabla_V(\varphi_*^{-1}(X)), V) \, d\mu \\ &= \int_G g(\nabla_{\varphi_*V} X, \varphi_*V) \, d\mu. \end{aligned}$$

□

For a function, we also have the following result.

**Lemma 4.2.** *With the notation above, for any function  $f$ ,*

$$\begin{aligned} \nabla \bar{f} &= \overline{\nabla f}, \\ \text{Hess } \bar{f} &= \int_G \text{Hess } f(\varphi_* V, \varphi_* V) \, d\mu. \end{aligned}$$

*Proof.* First, note that the second equation follows from the first combined with Lemma 4.1, along with the fact that

$$\text{Hess } f = \frac{1}{2} L_{\nabla f} g.$$

To prove the first equation, let  $V$  be a vector field on  $M$ , and observe that

$$\begin{aligned} g(\nabla \bar{f}, V) &= D_V \left( \int_G f \circ \varphi \, d\mu \right) = \int_G df(\varphi_* V) \, d\mu \\ &= \int_G g(\varphi_*^{-1}(\nabla f), V) = g(\overline{\nabla f}, V). \end{aligned} \quad \square$$

We are now ready to show that the weighted curvatures can be averaged over the compact group  $G$ . First, we consider the  $\infty$ -cases.

**Lemma 4.3.** *Given a triple  $(M, g, X)$  and a compact subgroup  $G$  of the isometry group, the weighted curvatures satisfy*

$$\begin{aligned} \text{Ric}_{\bar{X}}(U, V) &= \int_G \text{Ric}_X(\varphi_* U, \varphi_* V) \, d\mu, \\ \text{sec}_{\bar{X}}^V(U) &= \int_G \text{sec}_X^{\varphi_* V}(\varphi_* U) \, d\mu, \end{aligned}$$

where  $\bar{X}$  is the average of  $X$ . In particular, if  $\text{sec}_X \geq \lambda$ , then  $\text{sec}_{\bar{X}} \geq \lambda$  where  $\bar{X}$  is  $G$ -invariant.

**Remark 4.4.** One can similarly draw conclusions about upper bounds and for the Bakry-Emery Ricci curvature. In addition, analogous statements hold for  $\text{Ric}_f$  and  $\text{sec}_f$ . They follow immediately from Lemmas 4.2 and 4.3.

*Proof.* Using Lemma 4.1, we can see all we need to show is

$$\begin{aligned} \text{Ric}(U, V) &= \int_G \text{Ric}(\varphi_* U, \varphi_* V) \, d\mu \\ \text{sec}(U, V) &= \int_G \text{sec}(\varphi_* U, \varphi_* V) \, d\mu. \end{aligned}$$

But this just follows from the isometry invariance of the curvature as well as the fact that  $d\mu$  has unit volume. □

For the strongly weighted curvatures, averaging the vector field  $X$  causes some issues, as the equation contains terms that are quadratic in  $X$ . In the gradient case, we can overcome this by changing the form of the potential function. Given  $m$ , set  $u = e^{-f/m}$ ; then, a simple calculation shows that

$$\text{Hess } f - \frac{df \otimes df}{m} = -\frac{m \text{Hess } u}{u}.$$

So, we have

$$\text{Ric}_f^m = \text{Ric} - \frac{m \text{Hess } u}{u},$$

and, choosing  $m = -1$ ,

$$\overline{\text{sec}}_f^V(U) = \text{sec}(V, U) + \frac{\text{Hess } u}{u}(V, V).$$

In these cases, it is natural to average the function  $u$ .

Let  $\bar{u}(p) = \int_G u(\varphi(p)) \, d\mu$ , and define  $\bar{f} = -m \log(\bar{u})$ . Then, we have the following lemma.

**Lemma 4.5.** *Given a triple  $(M, g, f)$  and a compact subgroup  $G$  of the isometry group, the weighted curvatures satisfy*

$$\begin{aligned} \bar{u} \text{Ric}_{\bar{f}}^m(U, V) &= \int_G u \text{Ric}_f^m(\varphi_*U, \varphi_*V) \, d\mu, \\ \bar{u} \overline{\text{sec}}_{\bar{f}}^V(U) &= \int_G u \overline{\text{sec}}_f^{\varphi_*V}(\varphi_*U) \, d\mu, \end{aligned}$$

where  $\bar{u}$  is the average of  $u = e^{-f/m}$  and  $\bar{f} = -m \log(\bar{u})$ . In particular, if  $\overline{\text{sec}}_f \geq \lambda$ , then  $\overline{\text{sec}}_{\bar{f}} \geq \lambda$ , where  $\bar{f}$  is  $G$ -invariant.

*Proof.* We will discuss the Ricci case, and the sectional curvature case will follow from an analogous argument. We have

$$\begin{aligned} u \text{Ric}_f^m(V, V) &= u \text{Ric}(V, V) - m \text{Hess } u(V, V), \\ \int_G u \text{Ric}_f^m(\varphi_*V, \varphi_*V) \, d\mu &= \int_G (u \text{Ric}(\varphi_*V, \varphi_*V) - m \text{Hess } u(\varphi_*V, \varphi_*V)) \, d\mu \\ &= \bar{u} \text{Ric}(V, V) - m \text{Hess } \bar{u}(V, V) = \bar{u} \text{Ric}_{\bar{f}}^m(V, V). \end{aligned}$$

To see the final remark, note that, if  $\text{Ric}_f^m \geq \lambda g$ , then

$$\text{Ric}_{\bar{f}}^m(U, V) = \frac{\int_G u \text{Ric}_f^m(\varphi_*U, \varphi_*V) \, d\mu}{\bar{u}} \geq \frac{\int_G \lambda u g(U, V) \, d\mu}{\bar{u}} = \lambda g(U, V),$$

so  $\text{Ric}_{\bar{f}}^m \geq \lambda g$  as well. Similar arguments hold for upper bounds.  $\square$

We remind the reader that Lemmas 4.3 and 4.5 immediately imply Corollary 2.2: if  $(M, g, X)$  has positive weighted sectional curvature and  $G$  is a compact subgroup of the isometry group of  $(M, g)$ , then there exists a  $G$ -invariant vector field  $\bar{X}$  such that  $(M, g, \bar{X})$  has positive weighted sectional curvature. Indeed, if  $\text{sec}_X > 0$ , then one can replace  $X$  by its average  $\bar{X}$  over the  $G$ -orbits. If  $X = \nabla f$  and  $\overline{\text{sec}}_f > 0$ , then one can replace  $X$  by  $\bar{X} = \nabla \bar{f}$ , where  $\bar{f} = \log(\bar{u})$  and where  $\bar{u}$  is the average of  $u = e^f$  over the  $G$ -orbits.

**Remark 4.6.** Note that Lemma 4.5 does not clearly extend to the non-gradient case, since there is no globally defined function  $u$  to average. We can still average over  $X$ , but only one side of the curvature bound is preserved. To see this, note that the strongly weighted curvatures satisfy

$$\overline{\text{sec}}_X^V(U) = \int_G \overline{\text{sec}}_X^{\varphi_* V}(\varphi_* U) \, d\mu + \left( \int_G g(X, \varphi_* V) \, d\mu \right)^2 - \int_G g(X, \varphi_* V)^2 \, d\mu.$$

In particular, by the Cauchy-Schwarz inequality,

$$\overline{\text{sec}}_X^V(U) \leq \int_G \overline{\text{sec}}_X^{\varphi_* V}(\varphi_* U) \, d\mu,$$

so upper bounds on strongly weighted curvatures are preserved by averaging the density. Similar statements hold in the gradient case. For the  $m$ -Bakry-Emery curvature, we similarly have

$$\begin{aligned} \text{Ric}_X^m(V, V) &= \int_G \text{Ric}_X^m(\varphi_* V, \varphi_* V) \, d\mu \\ &\quad - \frac{1}{m} \left[ \left( \int_G g(X, \varphi_* V) \right)^2 - \int_G g(X, \varphi_* V)^2 \right]. \end{aligned}$$

**4.2. Homogeneous metrics.** We now apply averaging the density to the special case of homogeneous metrics. Homogeneous Riemannian manifolds with positive sectional curvature are classified Wallach [Wal72] and Bérard-Bergery [BB76]. By averaging the density, we show here that there are no additional examples in the weighted case when  $X = \nabla f$ .

**Proposition 4.7.** *Let  $(M, g)$  be a compact homogeneous manifold, and let also  $f \in C^\infty(M)$ . We have the following:*

- (1) *If  $\text{Ric}_f \geq \lambda g$  or  $\text{Ric}_f^m \geq \lambda g$ , then  $\text{Ric} \geq \lambda g$ .*
- (2) *If  $\text{sec}_f \geq \lambda g$  or  $\overline{\text{sec}}_f \geq \lambda g$ , then  $\text{sec} \geq \lambda g$ .*

*Analogous results hold for upper bounds.*

This proposition immediately implies Theorem B from the Introduction. Indeed, if  $(M, g)$  admits a gradient field  $X = \nabla f$  with positive weighted sectional curvature, then  $\overline{\text{sec}}_f > 0$ , and hence this proposition applies.

*Proof.* Let  $G$  be the isometry group of  $(M, g)$ . In all cases, we can replace  $f$  by a  $G$ -invariant function  $\tilde{f}$  such that the  $\tilde{f}$ -weighted curvatures have the same lower bounds as the  $f$ -weighted curvatures. Since  $G$  acts transitively,  $\tilde{f}$  is constant, so the  $\tilde{f}$ -weighted curvatures are equal to the usual, unweighted curvatures.  $\square$

It is not clear whether this fact is also true when the field  $X$  is not gradient. Averaging the field so that it is invariant under the isometries will not necessarily make the field Killing, but there is one important case where it does.

**Proposition 4.8.** *If a compact Lie group with a bi-invariant metric admits an  $X$  such that  $\text{sec}_X \geq \lambda$  or  $\text{Ric}_X \geq \lambda g$ , then  $\text{sec} \geq \lambda$  or  $\text{Ric} \geq \lambda g$ , respectively.*

*Proof.* We can replace  $X$  by its average over the left and right actions of  $G$ . This preserves the lower bounds on curvature, and makes  $X$  bi-invariant and hence a Killing field. Hence,  $L_X g = 0$ , so the weighted curvatures equal the unweighted curvatures.  $\square$

In particular, the previous two propositions have the following corollary.

**Corollary 4.9.** *A compact Lie group with a bi-invariant metric has positive weighted sectional curvature if and only if it has positive sectional curvature.*

In the simplest non-trivial case of a left-invariant metric not bi-invariant, a computation shows that we again do not get new examples.

**Proposition 4.10.** *If a left-invariant metric on the Lie group  $SU(2)$  supports a vector field  $X$  such that  $\text{sec}_X \geq \lambda$  or  $\text{Ric}_X \geq \lambda g$ , then  $\text{sec} \geq \lambda$  or  $\text{Ric} \geq \lambda g$ , respectively.*

*Proof.* For a left-invariant metric on  $SU(2)$ , choose an orthonormal frame  $\lambda_1^{-1}X_1, \lambda_2^{-1}X_2, \lambda_3^{-1}X_3$  such that  $[X_i, X_{i+1}] = 2X_{i+2}$  with indices taken mod 3. It follows that

$$\begin{aligned} \nabla_{X_i} X_i &= 0 \\ \nabla_{X_i} X_{i+1} &= \left( \frac{\lambda_{i+2}^2 + \lambda_{i+1}^2 - \lambda_i^2}{\lambda_{i+2}^2} \right) X_{i+2}, \\ \nabla_{X_{i+1}} X_i &= \left( \frac{-\lambda_{i+2}^2 + \lambda_{i+1}^2 - \lambda_i^2}{\lambda_{i+2}^2} \right) X_{i+2}. \end{aligned}$$

Now, since  $SU(2)$  is compact, we can assume by averaging that  $X$  is a left-invariant vector field, which we will write as  $X = a_1X_1 + a_2X_2 + a_3X_3$  for constants  $a_i$ . We have

$$\begin{aligned} (L_X g)(X_i, X_i) &= 2g(\nabla_{X_i} X, X_i) = 0, \\ (L_X g)(X_i, X_{i+1}) &= g(\nabla_{X_i} X, X_{i+1}) + g(\nabla_{X_{i+1}} X, X_i) \\ &= 2a_{i+2}(\lambda_i^2 - \lambda_{i+1}^2). \end{aligned}$$

This shows that  $X$  is not a Killing field in general. However,  $\sec_X(X_i, X_j) = \sec(X_i, X_j)$ , so if  $\sec_X \geq \lambda$ , then  $\sec(X_i, X_j) \geq \lambda$ . Further computation also shows that the basis  $X_1 \wedge X_2, X_2 \wedge X_3, X_3 \wedge X_1$  diagonalizes the curvature operator, and thus that all of the sectional curvatures are bounded by the maximum and minimum curvatures of the sectional curvatures involving  $X_1, X_2$ , and  $X_3$ . Thus, we actually have  $\sec \geq \lambda$ . The basis  $X_1, X_2, X_3$  also diagonalizes the Ricci tensor, so the statement about Ricci curvatures follows similarly.  $\square$

In general, Proposition 4.7 does not hold in the non-compact case, as we have already seen in Example 3.2. We can generalize the Gaussian example in the following simple way.

**Example 4.11.** Suppose that  $(M, g)$  is a simply connected space of non-positive sectional curvature. The distance function to a point squared,  $d^2$ , is a smooth function. Moreover,  $\text{Hess}(d^2) \geq 2g$ . Therefore, if  $(M, g)$  has sectional curvature bounded from below by  $-K$ , then, for  $f = Ad^2$ , we have  $\sec_f \geq 2A - K$ , which we can make arbitrarily large.

Letting  $(M, g)$  in the example be a hyperbolic space gives a noncompact homogeneous manifold with positive weighted sectional curvature and negative sectional curvature. We also note that there are many examples of non-compact homogeneous Ricci soliton metrics (i.e., metrics with  $\text{Ric}_X = \lambda g$ ) that do not have  $\text{Ric} = \lambda g$ . Examples of homogeneous metrics with  $\text{Ric}_f^m = \lambda g$  that do not have  $\text{Ric} = \lambda g$  are also constructed in [HPW15].

### 5. RIEMANNIAN SUBMERSIONS AND CHEEGER DEFORMATIONS

We analyze the behavior of the weighted and strongly weighted directional curvature operators under a Riemannian submersion  $\pi : M \rightarrow B$ . For this, we restrict ourselves to vector fields  $X$  on  $M$  for which the vector field  $\pi_*(X)$  on  $B$  is well defined. Following Besse [Bes08, Chapter 9], let  $R, \hat{R}$ , and  $\check{R}$  denote the curvature tensors of  $M$ , the fibers, and the base, respectively, and let  $\mathcal{V}$  and  $\mathcal{H}$  denote the projection maps onto the vertical and horizontal spaces, respectively.

**Theorem 5.1 (O’Neill formulas).** *Let  $(M, g)$  be a closed Riemannian manifold, let  $\pi$  be a Riemannian submersion with domain  $M$ , and let  $X$  be a smooth vector field on  $M$  such that the map  $p \mapsto \pi_*(X_p)$  is constant along the fibers of  $\pi$ . If  $Y$  and  $Z$  are horizontal vector fields and  $U$  and  $V$  are vertical vector fields on  $M$ , then*

$$\begin{aligned} R_X^V(U, U) &= \hat{R}_{VX}^V(U, U) + g(T_U V, T_U V) \\ &\quad - g(T_U U, T_V V) - g(T_V V, \mathcal{H}X)g(U, U), \\ R_X^Z(Y, Y) &= \check{R}_{\pi_* X}^{\pi_* Z}(\pi_* Y, \pi_* Y) - 3g(A_Y Z, A_Y Z), \end{aligned}$$

and likewise with  $R_X, \hat{R}_{VX}$  and  $\check{R}_{\pi_* X}$  replaced by the strongly weighted directional curvature operators on  $M$ , the fibers, and the base, respectively.

In particular, if  $(Y, Z)$  is an orthonormal pair of horizontal vector fields, then

$$\sec_{\pi_*X}^{\pi_*Y}(\pi_*Z) = \sec_X^Y(Z) + \frac{3}{4} |[Y, Z]^V|^2,$$

and likewise for  $\overline{\sec}_X$ .

Analogous statements hold in the gradient case. There, one assumes that  $f$  is a smooth function on  $M$  that is constant along the fibers of  $\pi$ . The function  $f$  replaces  $X$  in the above formulas, and the induced map  $\tilde{f}$  on the base replaces  $\pi_*X$ . The gradient case follows from the general case since  $d\tilde{f}$  and  $\text{Hess } \tilde{f}$  pull back via  $\pi$  to  $df$  and  $\text{Hess } f$ , respectively.

As for sectional curvature, the O'Neill formulas show that the base of a Riemannian submersion inherits lower bounds on weighted or strongly weighted sectional curvatures. In particular, if the total space admits a vector field  $X$  with positive weighted sectional curvature such that  $X$  descends to a well-defined vector field on the base, then the base too has positive weighted sectional curvature (see Corollary 2.3).

Finally, the vector field  $X$  is arbitrary, and hence need not be horizontal or vertical. For example, suppose  $\pi$  is the quotient map by a free, isometric group action. The vector field  $X$  might be an action field (hence, vertical), basic (hence, horizontal), or any smooth combination of the two (hence, neither).

*Proof.* Let  $\hat{g}$  and  $\check{g}$  denote the metrics on the fibers and the base, respectively. First, the conclusions in the strongly weighted cases follow immediately from the weighted cases, since

$$\begin{aligned} g(X, V)^2 g(U, U) &= \hat{g}(\mathcal{V}X, V)^2 \hat{g}(U, U), \\ g(X, Z)^2 g(Y, Y) &= \check{g}(\pi_*X, \pi_*Z)^2 \check{g}(\pi_*Y, \pi_*Y). \end{aligned}$$

Second, the weighted cases follow from the unweighted case once we establish that

$$\begin{aligned} \frac{1}{2} (L_X g)(V, V) g(U, U) &= \frac{1}{2} (L_{\mathcal{V}X} \hat{g})(V, V) \hat{g}(U, U) - g(T_V V, \mathcal{H}X) g(U, U), \\ \frac{1}{2} (L_X g)(Z, Z) g(Y, Y) &= \frac{1}{2} (L_{\pi_*X} \check{g})(\pi_*Z, \pi_*Z) \check{g}(\pi_*Y, \pi_*Y). \end{aligned}$$

Indeed these follow from the fact that  $U$  is vertical, the fact that  $Y$  is horizontal, and the observations that

$$\begin{aligned} \frac{1}{2} (L_X g)(V, V) &= g(\nabla_V X, V) \\ &= g(\nabla_V (\mathcal{V}X), V) + g(\nabla_V (\mathcal{H}X), V) \\ &= \hat{g}(\hat{\nabla}_V (\mathcal{V}X), V) - g(\nabla_V V, \mathcal{H}X) \\ &= \frac{1}{2} (L_{\mathcal{V}X} \hat{g})(V, V) - g(T_V V, \mathcal{H}X) \end{aligned}$$

and

$$\begin{aligned} \frac{1}{2}(L_X g)(Z, Z) &= g(\nabla_Z X, Z) = \check{g}(\nabla_{\pi_* Z} \pi_* X, \pi_* Z) \\ &= \frac{1}{2}(L_{\pi_* X} \check{g})(\pi_* Z, \pi_* Z). \end{aligned} \quad \square$$

Regarding the O’Neill formulas for mixed inputs (vertical and horizontal), we note that one simply obtains weighted versions by adding the appropriate terms from the definition of  $R_X$  and  $\check{R}_X$ . The formulas do not simplify as in Theorem 5.1, but one can still use them. To illustrate this with one easy example, we generalize here a result of Weinstein [Wei80, Theorem 6.1] to the case of positive weighted sectional curvature (cf. Florit and Ziller [FZ11] and Chen [Che14]).

**Theorem 5.2 (Weinstein).** *Let  $\pi : M \rightarrow B$  be a Riemannian submersion of closed Riemannian manifolds with totally geodesic fibers. If there then exists a function  $f \in C^\infty(M)$  such that  $\overline{\text{sec}}_f > 0$  on all orthonormal pairs of vectors spanning “vertical” planes, then the fiber dimension is at most  $\rho(\dim B)$ , where  $\rho(n)$  denotes the maximum number of linearly independent vector fields on  $\mathbb{S}^{n-1}$ .*

Note that this reduces to the Weinstein result when  $f = 0$ . Recall also that  $f \in C^\infty(M)$  is basic if it is constant along the fibers of  $\pi$ .

*Proof.* Since the fibers are totally geodesic, the  $T$  tensor vanishes. Hence, for any orthonormal pair  $(V, Z)$ , where  $V$  is vertical and  $Z$  is horizontal, the O’Neill formula

$$R(Z, V, V, Z) = |A_Z V|^2 - |T_Z U|^2 + g((\nabla_Z T)_V V, Z)$$

implies

$$\overline{\text{sec}}_f^V(Z) = |A_Z V|^2 + \text{Hess } f(V, V) + \text{d}f(V)^2.$$

At a point  $p \in M$  where  $f$  is maximized,  $\text{d}f(V) = 0$  and  $\text{Hess } f(V, V) \leq 0$  for all  $V$ . Hence,  $A_Z V \neq 0$  for all vertical pairs  $(V, Z)$  at  $p$ . The proof now proceeds as in [Wei80] by constructing  $\dim(\mathcal{V}_p)$  linearly independent vectors on the unit sphere in  $\mathcal{H}_p$ , where  $\mathcal{V}_p$  and  $\mathcal{H}_p$  are the vertical and horizontal spaces at  $p$ , respectively. □

Note that Theorem 5.2 relates to a conjecture of Fred Wilhelm, namely, that  $\dim(F) < \dim(B)$  for any Riemannian submersion from a manifold  $M$  with positive sectional curvature, where  $\dim(F)$  and  $\dim(B)$  denote the dimensions of the fibers and the base, respectively. If one only assumes  $\text{sec} > 0$  almost everywhere on  $M$ , then there are counterexamples due to Kerin [Ker11]. On the other hand, the above result suggests that the assumption of positive sectional curvature might be weakened to cover manifolds with density. For example, Frankel’s theorem (Theorem 7.1) in the weighted case implies the following: if  $M$  admits a vertical vector field  $X$  such that  $M$  has positive weighted sectional curvature, then the conclusion of Wilhelm’s conjecture holds.

As a second application, we discuss Cheeger deformations. These have been used in multiple constructions of metrics with positive or non-negative sectional curvature (see Ziller [Zil07] for a survey). Here, we establish the weighted curvature formulas for the deformed metric in terms of the original. We will use the formulas from this section in the proof of Theorem E.

The setup involves a Riemannian manifold  $(M, g)$ , a subgroup  $G$  of the isometry group, a bi-invariant metric  $Q$  on  $G$ , and a real parameter  $\lambda > 0$ . We are interested in understanding how the weighted curvatures behave under these perturbations. Hence, we also fix a smooth vector field  $X$  on  $M$ . We assume that  $X$  is  $G$ -invariant, which can be arranged if the subgroup  $G$  is compact (e.g., if  $G$  is closed and  $M$  is compact).

The new metric on  $M$  is denoted by  $g_\lambda$ . It is the metric for which the map  $\pi: (G \times M, \lambda Q + g) \rightarrow (M, g_\lambda)$  given by  $(h, p) \mapsto h^{-1}p$  is a Riemannian submersion. There is a  $(\lambda Q + g)$ -orthogonal decomposition of  $T_{(e,p)}(G \times M)$  as

$$\{(Y, Y_p^*) \mid Y \in \mathfrak{g}\} \oplus \{(-|Y_p^*|^2 Y, \lambda|Y|^2 Y_p^*) \mid Y \in \mathfrak{g}\} \oplus \{(0, Z) \mid Z \in T_p(G \cdot p)^\perp\}.$$

Here and throughout,  $\mathfrak{g} = T_e G$  denotes the Lie algebra of  $G$ , and  $Y^*$  denotes the Killing field associated with  $Y \in \mathfrak{g}$ . The first of these summands is the vertical space  $\mathcal{V}_{(p,e)} = \ker(D\pi_{(e,p)})$  of the projection  $\pi$ . The last two summands together form the horizontal space  $\mathcal{H}_{(e,p)} = \mathcal{V}_{(e,p)}^\perp$ .

The horizontal lift of  $Y^* \in T_p(G \cdot p) \subseteq T_p M$  is

$$\frac{1}{|Y_p^*|^2 + \lambda|Y|^2} (-|Y_p^*|^2 Y, \lambda|Y|^2 Y_p^*),$$

and the horizontal lift of  $Z \in T_p(G \cdot p)^\perp \subseteq T_p M$  is  $(0, Z)$ . Note that  $|Z|_{g_\lambda} = |Z|_g$ , while

$$|Y^*|_{g_\lambda}^2 = \frac{\lambda|Y|^2 |Y^*|^2}{|Y_p^*|^2 + \lambda|Y|^2}.$$

As  $\lambda \rightarrow \infty$ ,  $|Y^*|_{g_\lambda}$  increases and converges to  $|Y^*|_g$ , and hence  $|Y^*|_{g_\lambda} \leq |Y^*|$ . We will use this in the proof of the connectedness lemma.

Our goal now is to compute the weighted and strongly weighted directional curvature operators of  $(M, g_\lambda, X)$  in terms of those of  $(M, g, X)$ .

**Lemma 5.3 (Curvature tensors after Cheeger deformations).** *Let  $R = R^g$  and  $R^{g_\lambda}$  denote the curvature tensors of  $(M, g)$  and  $(M, g_\lambda)$ , respectively. For vector fields  $W_i$  on  $M$ , if  $\tilde{W}_i = (\tilde{W}_i^G, \tilde{W}_i^M)$  denote the horizontal lifts in  $G \times M$ , then*

$$\begin{aligned} g_\lambda((R^{g_\lambda})_X^{W_1}(W_2), W_3) &= \lambda Q((R^Q)^{\tilde{W}_1^G}(\tilde{W}_2^G), \tilde{W}_3^G) \\ &\quad + g((R^g)_X^{\tilde{W}_1^M}(\tilde{W}_2^M), \tilde{W}_3^M) \\ &\quad + (\lambda Q + g)(A_{\tilde{W}_1} \tilde{W}_2, A_{\tilde{W}_1} \tilde{W}_3). \end{aligned}$$

In particular, if  $Z_1$  and  $Z_2$  are vector fields in  $M$  that are everywhere orthogonal to the  $G$ -orbits, then

$$g_\lambda((R^{g_\lambda})_X^{Z_1}(Z_2), Z_2) \geq g((R^g)_X^{Z_1}(Z_2), Z_2).$$

If, in addition,  $(Z_1, Z_2)$ , forms an orthonormal pair with respect to  $g$  (equivalently, with respect to  $g_\lambda$ ), then  $(\sec^{g_\lambda})_X^{Z_1}(Z_2) \geq (\sec^g)_X^{Z_1}(Z_2)$ .

*Proof.* Consider the vector field  $(0, X)$  on  $G \times M$ . It is  $G$ -invariant, and  $\pi_*(0, X) = X$ , where  $\pi : (G \times M, \lambda Q + g) \rightarrow (M, g_\lambda)$  is the Riemannian submersion defining  $g_\lambda$ . The first claim follows directly from the (first) O’Neill formula in the weighted case applied to  $\pi$ . The second and third claims follow from the fact that the horizontal lift of  $Z \in T_p(G \cdot p)^\perp$  is  $(0, Z) \in T(G \times M)$ .  $\square$

### 6. WEINSTEIN’S FIXED POINT THEOREM AND APPLICATIONS

In the next two sections, we demonstrate how Synge-type arguments extend to the case of positive weighted sectional curvature. The only technical ingredient required is Lemma 2.4. We first prove Weinstein’s fixed point theorem in the weighted case.

**Theorem 6.1 (Weinstein’s fixed point theorem).** *Let  $(M^n, g)$  be a closed, orientable Riemannian manifold equipped with vector field  $X$  such that  $(M, g, X)$  has positive weighted sectional curvature. If  $F$  is an isometry of  $M$  with no fixed point, then  $F$  reverses orientation if  $n$  is even, and preserves it if  $n$  is odd.*

*Proof.* Corollary 2.2 implies we may assume without loss of generality that  $X$  is invariant under isometries. In particular,  $F_*(X) = X$ .

The proof now proceeds as in Weinstein [Wei68]. Using compactness, choose  $p \in M$  such that  $d(p, F(p))$  is minimal. Choose a unit-speed, minimizing geodesic  $\gamma : [a, b] \rightarrow M$  from  $p$  to  $F(p)$ . As in [Wei68], there exists a special unit-length, parallel vector field  $V$  along  $\gamma$ , and it suffices to show that the index  $I(V, V)$  of  $\gamma$  is negative. One of the properties of  $\gamma$  is that  $F_*(\gamma'(a)) = \gamma'(b)$ . By Lemma 2.4, it suffices to show that

$$g(\gamma'(t), X_{\gamma(t)}) \Big|_{t=a}^{t=b} = \langle \gamma'(b), X_{\gamma(b)} \rangle - \langle \gamma'(a), X_{\gamma(a)} \rangle = 0.$$

Indeed, this is the case since  $F$  carries  $\gamma'(a)$  to  $\gamma'(b)$  and  $X_{\gamma(a)}$  to  $X_{F(\gamma(a))} = X_{\gamma(b)}$ .  $\square$

We derive three corollaries of Weinstein’s theorem, all of which are analogues of what happens in the unweighted case. The first is the textbook application of Weinstein’s theorem to prove Synge’s theorem.

**Corollary 6.2 (Synge’s theorem).** *If  $(M^n, g, X)$  is closed and also has positive weighted sectional curvature, then the following hold:*

- If  $n$  is odd, then  $M$  is orientable.
- If  $n$  is even and  $M$  is orientable, then  $\pi_1(M)$  is trivial.

This is proved in [Wy115], but we indicate another proof based on Weinstein's theorem. Depending on whether  $n$  is odd or even, one applies Weinstein's theorem in the weighted case to the free action of  $\mathbb{Z}_2$  or  $\pi_1(M)$ , respectively, on the orientation or universal cover of  $M$  equipped with the pullback metric and vector field or function. For this, it is important that  $\pi_1(M)$  is finite (see Theorem 2.1).

Weinstein's theorem, together with O'Neill's formula, also provides another proof of Berger's result (see [BER66, GS94]).

**Corollary 6.3 (Berger's theorem).** *If  $(M^n, g, X)$  is closed and has positive weighted sectional curvature, then the following hold:*

- *If  $n$  is even, then any Killing field has a zero. Equivalently, any isometric torus action has a fixed point.*
- *If  $n$  is odd, any torus acting isometrically on  $M$  has a circle orbit. In particular, there exists a codimension-one subtorus that has a fixed point.*

We remark that the even-dimensional case is also proved in [Wy115].

*Proof.* The equivalence of the conclusions about Killing fields and torus actions is based on the fact that the isometry group of  $M$  is a compact Lie group. Consider an isometric action on  $M$  by a torus  $T$ . Without loss of generality, we assume that  $X$  is invariant under the action of  $T$ . The conclusion follows by choosing  $F \in T$  that generates a dense subgroup of  $T$ , and applying Weinstein's theorem to  $F$ .

The odd-dimensional case follows from the even-dimensional case and the O'Neill formula, as proved in Grove and Searle [GS94]). Since the even-dimensional case and O'Neill's formula hold in the weighted case, the proof is complete.  $\square$

Finally, it was observed in [Ken] that Weinstein's theorem pairs nicely with a result of Davis and Weinberger to provide an obstruction to free group actions on positively curved rational homology spheres of dimension  $4k + 1$ , as follows.

**Theorem 6.4 (Davis-Weinberger factorization).** *Let  $(M^{4k+1}, g, X)$  be closed with positive weighted sectional curvature. If the universal cover of  $M$  is a rational homology sphere, then  $\pi_1(M) \cong \mathbb{Z}_{2^e} \times \Gamma$  for some odd-order group  $\Gamma$ .*

*Proof.* Since  $\pi_1(M)$  is finite (see Theorem 2.1), we may consider the free action of  $\pi_1(M)$  on the universal cover of  $M$ , which is a compact, simply connected manifold with the same weighted curvature bound as  $M$ . By Weinstein's theorem in the weighted case, the action of  $\pi_1(M)$  is (rationally) homologically trivial. As  $\dim(M) \equiv 1 \pmod{4}$  and the surgery semicharacteristic  $\sum_{i \leq 2k} (-1)^i \dim H^i(M; \mathbb{Q})$  is odd, the factorization of  $\pi_1(M)$  follows from Theorem D in [Dav83].  $\square$

## 7. FRANKEL'S THEOREM AND WILKING'S CONNECTEDNESS LEMMA

In this section, we prove generalizations of Frankel's theorem and Wilking's connectedness lemma in the weighted case. Specifically, we assume throughout this

section that  $(M^n, g, X)$  is a Riemannian manifold equipped with a vector field  $X$  such that  $(M, g, X)$  has positive weighted sectional curvature.

**Theorem 7.1 (Frankel).** *Assume  $(M^n, g, X)$  is closed with positive weighted sectional curvature. Assume  $N_1$  and  $N_2$  are closed, totally geodesic submanifolds of  $M$  such that  $X$  is tangent to  $N_i$  for  $i \in \{1, 2\}$ . If  $\dim(N_1) + \dim(N_2) \geq n$ , then  $N_1$  and  $N_2$  intersect.*

Before proving this, we record an easy corollary we will use in the next section.

**Corollary 7.2.** *Let  $(M^n, g, X)$  be closed with positive weighted sectional curvature. Suppose  $G_1$  and  $G_2$  are subgroups of the isometry group of  $M$ , and suppose that  $N_1$  and  $N_2$  are components of the fixed-point sets of  $G_1$  and  $G_2$ , respectively. If  $\dim(N_1) + \dim(N_2) \geq n$ , then the submanifolds intersect.*

To deduce the corollary, one replaces  $X$  by  $\tilde{X}$  such that  $\tilde{X}$  is invariant under isometries of  $(M, g)$  and  $(M, g, \tilde{X})$  has positive weighted sectional curvature (see Corollary 2.2). For  $p \in N_1$ , it follows that  $X_p \in (T_p M)^{G_1} = T_p N_1$ ; hence,  $X$  is tangent to  $N_1$ , and likewise for  $N_2$ . The corollary follows since the  $N_i$  are closed and totally geodesic.

**Remark 7.3.** Note that both Theorem 7.1 and the corollary fail if we remove the assumption that  $M$  is compact. Indeed, consider the flat metric on Euclidean space, and let  $f = \frac{1}{2}d^2$ , where  $d$  is the distance to a fixed point in  $M$ . Clearly,  $\text{sec}_f^V(U) = \text{Hess } f(V, V) = 1$  for all orthonormal pairs  $(U, V)$ , yet any two parallel hyperplanes are disjoint, closed, totally geodesic, and have dimensions adding up to at least  $\dim M$ .

In fact,  $N_1$  and  $N_2$  are fixed-point components of reflection subgroups  $G_1 \cong \mathbb{Z}_2$  and  $G_2 \cong \mathbb{Z}_2$  of the isometry group. However, the subgroup generated by  $G_1$  and  $G_2$  is infinite, so we cannot replace  $X$  by a  $G_1$ - and  $G_2$ -invariant vector field as in Corollary 2.2 and proceed as in the proof of the corollary.

*Proof of Frankel’s theorem.* Let  $M, N_1, N_2$ , and  $X$  be as in the theorem. We proceed now as in Frankel [Fra61]. By compactness, there is a minimizing geodesic  $\gamma : [a, b] \rightarrow M$  connecting  $N_1$  to  $N_2$ . By the first variation formula,  $\gamma$  is normal to  $N_1$  and  $N_2$  at its endpoints. Since  $X$  is tangent to  $N_1$  and  $N_2$ ,

$$g(\gamma'(b), X_{\gamma(b)}) = g(\gamma'(a), X_{\gamma(a)}) = 0.$$

Using Lemma 2.4, the rest of the proof is as in the unweighted case. □

Wilking proved a vast generalization of Frankel’s result (see Theorem 2.1 in [Wil03]). The generalization to the weighted case is the following result.

**Theorem 7.4 (Wilking’s connectedness lemma).** *Let  $(M^n, g, X)$  be closed with positive weighted sectional curvature. The following hold:*

- (1) *If  $X$  is tangent to  $N^{n-k}$ , a closed, totally geodesic, embedded submanifold of  $M$ , then the inclusion  $N \rightarrow M$  is  $(n - 2k + 1)$ -connected.*

- (2) If  $X$  and  $N^{n-k}$  are as above, and if  $G$  acts isometrically on  $M$ , fixes  $N$  pointwise, and has principal orbits of dimension  $\delta$ , then the inclusion  $N \rightarrow M$  is  $(n - 2k + 1 + \delta)$ -connected.
- (3) If  $X$  is tangent to  $N_1^{n-k_1}$  and  $N_2^{n-k_2}$ , a pair of closed, totally geodesic, embedded submanifolds with  $k_1 \leq k_2$ , then  $N_1 \cap N_2 \rightarrow N_2$  is  $(n - k_1 - k_2)$ -connected.

As in the corollary to Frankel’s theorem, this result applies to inclusions of fixed-point components of isometric group actions.

*Proof.* The proof in each case proceeds as in Wilking [Wil03, Theorem 2.1], where the result is reduced to an index estimate. In the first and third cases, this estimate involves parallel vector fields, and hence extends to the weighted case exactly as in the proof of Frankel’s theorem above in the weighted case.

In the remaining case, the index estimate is a bit more involved, so we repeat it here, modifying it as necessary to cover the weighted case. The setup in [Wil03] is as follows. The metric  $g_\lambda$  on  $M$  is a Cheeger deformation of  $g$ , there is a geodesic  $c : [a, b] \rightarrow M$  that starts and ends perpendicular to  $N$ , and there is a  $(n - 2k + 1 + \delta)$ -dimensional vector space  $W$  of vector fields  $V$  along  $c$  such that the following hold:

- $V$  is tangent to  $N$  at the endpoints of  $c$ .
- $V$  is orthogonal to the  $G$ -orbits at all points along  $c$ .
- $V' = \nabla_{c'} V$  is tangent to the  $G$ -orbits at all points.

By the argument in [Wil03], it suffices to show that, for all  $V \in W$ , there exists  $\lambda > 0$  such that the index form with respect to  $g_\lambda$  of  $c$  evaluated on  $V$  is negative. We show this first under the assumption that  $\text{sec}_X > 0$  on  $M$ .

By Equation 2.1, the index form can be written as

$$\int_a^b (|V'|_{g_\lambda}^2 - (R^{g_\lambda})_{c'}^X(V, V) - 2g_\lambda(c', X)g_\lambda(V, V')) dt + g_\lambda(c', X) |V|_{g_\lambda}^2 \Big|_{t=a}^{t=b}.$$

First, we show that the last term in this expression is zero. Without loss of generality, we may assume that  $X$  is  $G$ -invariant and hence tangent to  $N$ . Since the  $G$ -orbits in  $N$  are trivial,  $X$  is orthogonal to the orbits. Hence, the horizontal lift of  $X_{c(t)}$  at  $t \in \{a, b\}$  is  $(0, X_{c(t)})$ , and

$$g_\lambda(c', X) \Big|_{t=a}^{t=b} = g(c', X) \Big|_{t=a}^{t=b} = 0.$$

Second, the O’Neill formula in the weighted case implies  $(R^{g_\lambda})_{c'}^X(V, V) \geq R_X^c(V, V)$ . Since this lower bound is independent of  $\lambda > 0$ , the proof will be complete once we show both of the following:

- $|V'|_{g_\lambda}^2 \rightarrow 0$  as  $\lambda \rightarrow 0$ .
- $g_\lambda(c', X)g_\lambda(V, V') \rightarrow 0$  as  $\lambda \rightarrow 0$ .

Indeed, since  $V'$  is tangent to the  $G$ -orbits,  $|V'|_{g_\lambda} \rightarrow 0$  as  $\lambda \rightarrow 0$ . This proves the first statement. The second statement follows from the first, together with the estimate

$$|g_\lambda(c', X)| |g_\lambda(V, V')| \leq |c'|_{g_\lambda} |X|_{g_\lambda} |V|_{g_\lambda} |V'|_{g_\lambda} \leq |c'|_g |X|_g |V|_g |V'|_{g_\lambda}.$$

Here, the second inequality follows since Cheeger deformations (weakly) decrease lengths, that is,  $|\cdot|_{g_\lambda} \leq |\cdot|_g$  for all  $\lambda > 0$ .

This completes the proof if  $\text{sec}_X > 0$ . Consider now the case where  $X = \nabla f$  and  $\overline{\text{sec}}_f > 0$ . Here, we consider the vector space

$$W_f = \{Y = e^f V \mid V \in W\},$$

and show that, for all  $Y \in W_f$ , there exists  $\lambda > 0$  such that the index  $I_c(Y, Y)$  of  $Y$  along  $c$  is negative. Since  $\dim(W_f) = \dim(W)$ , this would complete the proof in this case. This is easily accomplished by proceeding as in the previous case and using the alternative formula for the index given in Equation 2.2.  $\square$

### 8. TORUS ACTIONS AND POSITIVE WEIGHTED SECTIONAL CURVATURE

Throughout this section, we consider closed Riemannian manifolds  $(M, g)$  that are equipped with a vector field  $X$  such that  $(M, g, X)$  has positive weighted sectional curvature. In addition, we assume a torus  $T$  acts isometrically on  $M$ . Applying Corollary 2.2, if necessary, we assume  $X$  is invariant under the torus action.

Our first result is the following generalization of a result of Grove-Searle [GS94].

**Theorem 8.1 (Maximal symmetry rank).** *Let  $(M^n, g, X)$  be closed with positive weighted sectional curvature. If  $T^r$  is a torus acting effectively by isometries on  $M$ , then  $r \leq \lfloor (n + 1)/2 \rfloor$ . Moreover, if equality holds and  $M$  is simply connected, then  $M$  is homeomorphic to  $S^n$  or  $\mathbb{C}\mathbb{P}^{n/2}$ .*

The upper bound on  $r$  is sharp and agrees with Grove and Searle’s result. However, in the unweighted case, Grove and Searle prove an equivariant diffeomorphism classification when the maximal symmetry rank is achieved. We obtain this weaker rigidity statement by a different argument that relies on Wilking’s connectedness lemma and a lemma in Fang and Rong [FR05]. For a more detailed argument along these lines, we refer to [Pet06, Section 7.1.3].

*Proof.* By Berger’s theorem (Corollary 6.3) in the weighted case, there exists  $x \in M$  fixed by either  $T^r$  or a subtorus  $T^{r-1}$ , according to whether  $n$  is even or odd. Since this subtorus embeds into  $\text{SO}(n)$  via the isotropy representation, it follows that  $r \leq \lfloor (n + 1)/2 \rfloor$ .

We proceed to the equality case. First, if  $n \in \{2, 3\}$ , then  $M$  is homeomorphic to a sphere since it is simply connected by the resolution of the Poincaré

conjecture. Suppose therefore that  $n \geq 4$ . By arguing inductively as in Grove-Searle, it follows that some circle in  $T^r$  fixes a codimension-two submanifold  $N$ . By the connectedness lemma in the weighted case, we conclude that the inclusion  $N \hookrightarrow M$  is  $\dim(N)$ -connected. It follows immediately from the Poincaré duality that  $M$  and  $N$  are integral cohomology spheres or complex projective spaces (see, e.g., [Wil03, Section 7]). If  $M$  is an integral sphere, then it is a homeomorphism sphere by the resolution of the Poincaré conjecture. If  $M$  is an integral complex projective space, then the fact that  $N$  represents the generator of  $H^2(M; \mathbb{Z})$  implies that  $M$  is homeomorphic to complex projective space, by Lemma 3.6 in Fang-Rong [FR05].  $\square$

We note that there are a number of generalizations of Grove and Searle's result. These include results of Rong and Fang in the cases of "almost maximal symmetry rank" or non-negative curvature (see Fang and Rong [Ron02, FR05], Galaz-Garcia and Searle [GGS11, GGS14], and Wiemeler [Wie15]).

Returning to the case of positive curvature, there are additional results that assume less symmetry. We focus here on the following homotopy classification due to Wilking [Wil03, Theorem 2].

**Theorem 8.2 (Wilking's homotopy classification).** *Let  $M^n$  be a closed, simply connected, positively curved manifold, and let  $T^r$  act effectively by isometries on  $M$ . If  $n \geq 10$  and  $r \geq n/4 + 1$ ; then,  $M$  is either homeomorphic to  $S^n$  or  $\mathbb{H}\mathbb{P}^{n/4}$  or homotopy equivalent to  $\mathbb{C}\mathbb{P}^{n/2}$ .*

By Grove and Searle [GS94] and Fang and Rong [FR05], this result actually holds for all  $n \neq 7$ . Additionally, the conclusion in this theorem has been improved to a classification up to tangential homotopy equivalence (see Dessai and Wilking [DW04, Remark 1.4]). We prove the following analogue of Wilking's classification under a slightly stronger symmetry assumption.

**Theorem 8.3.** *Let  $(M^n, g, X)$  be closed and simply connected, and with positive weighted sectional curvature. If  $M$  admits an effective, isometric torus action of rank  $r \geq n/4 + \log_2 n$ , then  $M$  is homeomorphic to  $S^n$  or tangentially homotopy equivalent to  $\mathbb{C}\mathbb{P}^{n/2}$ .*

Note that  $\mathbb{H}\mathbb{P}^{n/4}$  does not appear in the conclusion. This is consistent with Theorem 3 in Wilking [Wil03], which states that the maximal rank of a smooth torus action on an integral  $\mathbb{H}\mathbb{P}^m$  is  $m + 1$ .

One reason for the larger symmetry assumption is that Wilking's original proof invokes the full strength of Grove and Searle's equivariant diffeomorphism classification. Since we do not prove this here, we cannot use exactly the same proof. In addition, the larger symmetry assumption allows us to side-step some of the more delicate parts of Wilking's proof, and thereby allows for a quick argument that captures the essence of his induction machinery, as described in the introduction of [Wil03].

*Proof of Theorem 8.3.* We first note that it suffices to prove that  $M$  has the integral cohomology of  $S^n$  or  $\mathbb{C}\mathbb{P}^{n/2}$ . Indeed, a simply connected integral sphere

is homeomorphic to the standard sphere by the resolution of the Poincaré conjecture. Moreover, it is well known that a simply connected integral complex projective space is homotopy equivalent to the standard one, and the classification up to tangential homotopy follows directly from Dessai and Wilking [DW04].

Second, note that the theorem holds in dimensions  $n \leq 13$  by the extension of Grove and Searle's result (Theorem 8.1). We proceed by induction for dimensions  $n \geq 14$ . By examining the isotropy representation at a fixed-point of  $T^r$  (or  $T^{r-1}$  in the odd-dimensional case), one sees that an involution  $\iota \in T^r$  exists such that some component  $N$  of its fixed-point set has codimension  $\text{cod}(N) \leq (n+3)/4$  (see, e.g., [Ken14, Lemma 1.8.(1)]). By replacing  $\iota$  by another involution, if necessary, we may assume  $\text{cod}(N)$  is minimal. In particular, the induced action of the torus  $T^r / \ker(T^r|_N)$  has rank at least  $r-1$ .

If  $\text{cod}(N) = 2$  and  $N$  is fixed by a circle, the claim follows as in the proof of the Grove-Searle result. Otherwise,  $T^r / \ker(T^r|_N)$  is a torus that acts effectively and isometrically on  $N$  with dimension at least  $\frac{1}{4} \dim N + \log_2(\dim N)$ . Since  $N$  is a fixed-point set of an involution in  $T^r$ , the vector field  $X$  is tangent to  $N$ , and  $N$  inherits positive weighted sectional curvature. By the connectedness lemma,  $N$  is simply connected. By the induction hypothesis,  $N$  is an integral sphere or complex projective space. By the connectedness lemma again, it follows that  $M$  too is an integral sphere or projective space. This concludes the proof.  $\square$

The theorems of this section should be viewed as a representative, as opposed to exhaustive, list of the kinds of topological results we can now generalize to the weighted setting. Indeed, the tools discussed in this paper have been applied to similar, weaker topological classification problems for positively curved manifolds with torus symmetry. Invariants calculated or estimated include the fundamental group (see Wilking [Wil03, Theorem 4], Frank-Rong-Wang [FRW13], Sun-Wang [SW09], and [Ken]), the Euler characteristic (see work of the first author and Amann [Ken13, AK14, AK]), and the elliptic genus (see Dessai [Des05, Des07] and Weisskopf [Wei]). Much of this work now can also be extended to the weighted case by using the results in this article.

On the other hand, it is much less clear whether some other prominent classification theorems for manifolds with positive curvature and torus symmetry can be extended to the weighted setting. Principal among these is the situation in low dimensions. In Section 3, we discussed why closed manifolds with positive weighted sectional curvature in dimension two and three are diffeomorphic to spherical space forms. In dimension four, Hsiang and Kleiner [HK89] proved that a closed, simply connected manifold  $M$  with positive curvature and an isometric circle action is homeomorphic to  $\mathbb{S}^4$  or  $\mathbb{C}\mathbb{P}^2$ . This result has been generalized in a number of ways. Recently, Grove and Wilking strengthened the conclusion to state that the circle action on  $M$  is equivariantly diffeomorphic to a linear action on one of these two spaces (see [GW14] and references therein for a survey of related work). A natural question is whether this result also holds for positive weighted sectional curvature.

**Question 8.4.** *Let  $(M^4, g, X)$  be simply connected and closed, and with positive weighted sectional curvature. Is every effective, isometric circle action on  $M$  equivariantly diffeomorphic to a linear action on  $S^4$  or  $\mathbb{C}P^2$ ?*

In dimension five, Rong [Ron02] proved that a positively curved  $M^5$  with an isometric 2-torus action is diffeomorphic to  $S^5$ . This result has also been improved to an equivariant diffeomorphism classification (see Galaz-Garcia and Searle [GGS14]), giving the following question.

**Question 8.5.** *Let  $(M^5, g, X)$  be simply connected and closed, and with positive weighted sectional curvature. Is every effective, isometric torus action of rank two on  $M$  equivariantly diffeomorphic to a linear action on  $S^5$ ?*

## 9. FUTURE DIRECTIONS

In addition to addressing Questions 8.4 and 8.5, another avenue of research is to consider compact manifolds with density that admit positive weighted curvature and an isometric action by an arbitrary Lie group  $G$ . To make this problem tractable, one can assume that  $G$  is large in some sense, for example, that  $G$  or its principal orbits have large dimension. Notable are classification results in this context due to Wallach [Wal72] and Bérard-Bergery [BB76] for transitive group actions, Wilking [Wil06] for more general group actions, Grove and Searle [GS97] and Spindeler [Spi14] for fixed-point homogeneous group actions, and Grove and Kim [GK04] for fixed-point cohomogeneity-one group actions. In the non-negatively curved case, especially in small dimensions, there have been some extensions of these results due to DeVito [DeV14, DeV], Galaz-Garcia and Spindeler [GGS12], Simas [Sim], and Gozzi [Goz15].

A particularly interesting case is where  $G$  is so large that the principal orbits have codimension one. Manifolds that admit a cohomogeneity-one metric with positive sectional curvature have been classified by Verdiani [Ver04] in the even-dimensional case, and by Grove, Wilking, and Ziller [GWZ08] in the odd-dimensional case (see also [VZ14] and the recent generalization to the case of polar actions by Fang, Grove, and Thorbergsson [FGT]).

The classification is actually incomplete in dimension seven, as there are two infinite families of manifolds that are considered “candidates” to admit positive curvature. There are very few examples of manifolds that admit positive curvature, so it was remarkable that one of these candidates was recently shown to admit positive sectional curvature by Darricott [Dea11] and Grove-Verdiani-Ziller [GVZ11]. It remains to be seen whether the others admit positive curvature.

It would be interesting to examine these results in the case of manifolds with density. Doing this would hopefully lead to new insights into the question posed in the Introduction: if  $(M, g, X)$  is compact with positive weighted sectional curvature, does  $M$  admit a metric with positive sectional curvature?

The most prominent missing ingredient when trying to generalize results to the weighted setting is a Toponogov-type triangle comparison theorem and the resulting convexity properties of distance functions. These crucial tools would be

needed to address Questions 8.4 and 8.5, the equivariant diffeomorphism rigidity in Grove and Searle's theorem (Theorem 8.1), and the results above for general group actions.

The examples in Section 3 show that the classical statement of the Toponogov theorem is false for positive weighted sectional curvature. On the other hand, we can make an analogy here with the situation of Ricci curvature and Bakry-Emery Ricci curvature. For positive Ricci curvature, instead of convexity of the distance function, one obtains Laplacian and volume comparisons. These comparisons, strictly speaking, do not hold for positive Bakry-Emery Ricci curvature, but they have modified weaker versions that are still enough to recover topological obstructions (see [WW09]). We believe there should be some form of modified convexity for distance functions one obtains from positive weighted sectional curvature, which may lead to generalizations of all of the results mentioned above. This will be the topic of future research.

**Acknowledgements.** We would like to thank Karsten Grove, Frank Morgan, Guofang Wei, Dmytro Yeroshkin, and Wolfgang Ziller for helpful suggestions and discussions. The first author is partially supported by NSF grants (nos. DMS-1045292 and DMS-1404670). The second author is partially supported by a grant from the Simons Foundation (no. 355608, William Wylie).

## REFERENCES

- [AK] M. AMANN AND L. KENNARD, *On a generalized conjecture of Hopf with symmetry*, Compos. Math. **153** (2017), no. 2, 313–322. <http://dx.doi.org/10.1112/S0010437X16008150>. MR3604865.
- [AK14] ———, *Topological properties of positively curved manifolds with symmetry*, Geom. Funct. Anal. **24** (2014), no. 5, 1377–1405. <http://dx.doi.org/10.1007/s00039-014-0300-9>. MR3261629.
- [BB76] L. BERARD-BERGERY, *Les variétés riemanniennes homogènes simplement connexes de dimension impaire à courbure strictement positive*, J. Math. Pures Appl. (9) **55** (1976), no. 1, 47–67 (French). MR0417987.
- [BÉ85] D. BAKRY AND M. ÉMERY, *Di usions hypercontractives*, Séminaire de probabilités, XIX, 1983/84, Lecture Notes in Math., vol. 1123, Springer, Berlin, 1985, pp. 177–206 (French). <http://dx.doi.org/10.1007/BFb0075847>. MR889476.
- [BER66] M. BERGER, *Trois remarques sur les variétés riemanniennes à courbure positive*, C. R. Acad. Sci. Paris Sér. A-B **263** (1966), A76–A78 (French). MR0199823.
- [Bes08] A. L. BESSE, *Einstein Manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition. MR2371700.
- [CG72] J. CHEEGER AND D. GROMOLL, *On the structure of complete manifolds of nonnegative curvature*, Ann. of Math. (2) **96** (1972), no. 3, 413–443. <http://dx.doi.org/10.2307/1970819>. MR0309010.
- [Che14] X. CHEN, *Curvature and Riemannian Submersions*, Ph.D. Thesis (2014), University of Notre Dame, Ann Arbor, MI.
- [CHH<sup>+</sup>06] I. CORWIN, N. HOFFMAN, S. HURDER, V. SESUM, AND Y. XU, *Differential geometry of manifolds with density*, Rose-Hulman Und. Math. J. **7** (2006), no. 1.
- [CLT06] X. CHEN, P. LU, AND G. TIAN, *A note on uniformization of Riemann surfaces by Ricci flow*, Proc. Amer. Math. Soc. **134** (2006), no. 11, 3391–3393 (electronic). <http://dx.doi.org/10.1090/S0002-9939-06-08360-2>. MR2231924.

- [CM11] I. CORWIN AND F. MORGAN, *The Gauss-Bonnet formula on surfaces with densities*, *Involvement* **4** (2011), no. 2, 199–202. <http://dx.doi.org/10.2140/involve.2011.4.199>. MR2876199.
- [Dav83] J. F. DAVIS, *The surgery semicharacteristic*, *Proc. London Math. Soc.* (3) **47** (1983), no. 3, 411–428. <http://dx.doi.org/10.1112/plms/s3-47.3.411>. MR716796.
- [Dea11] O. DEARRICOTT, *A 7-manifold with positive curvature*, *Duke Math. J.* **158** (2011), no. 2, 307–346. <http://dx.doi.org/10.1215/00127094-1334022>. MR2805071.
- [Des05] A. DESSAI, *Characteristic numbers of positively curved Spin-manifolds with symmetry*, *Proc. Amer. Math. Soc.* **133** (2005), no. 12, 3657–3661 (electronic). <http://dx.doi.org/10.1090/S0002-9939-05-07928-1>. MR2163604.
- [Des07] ———, *Obstructions to positive curvature and symmetry*, *Adv. Math.* **210** (2007), no. 2, 560–577. <http://dx.doi.org/10.1016/j.aim.2006.07.003>. MR2303232.
- [DeV14] J. DEVITO, *The classification of compact simply connected biquotients in dimensions 4 and 5*, *Differential Geom. Appl.* **34** (2014), 128–138. <http://dx.doi.org/10.1016/j.difgeo.2014.04.002>. MR3209541.
- [DeV] ———, *The classification of compact simply connected biquotients in dimensions 6 and 7*, *Math. Ann.*, available at <https://arxiv.org/abs/1403.6087>.
- [DW04] A. DESSAI AND B. WILKING, *Torus actions on homotopy complex projective spaces*, *Math. Z.* **247** (2004), no. 3, 505–511. <http://dx.doi.org/10.1007/s00209-003-0614-z>. MR2114425.
- [FGT] F. FANG, K. GROVE, AND G. THORBERGSSON, *Tits geometry and positive curvature*, *Acta Math.*, to appear, available at <http://arxiv.org/abs/arXiv:1205.6222>.
- [FR05] F. FANG AND X. RONG, *Homeomorphism classification of positively curved manifolds with almost maximal symmetry rank*, *Math. Ann.* **332** (2005), no. 1, 81–101. <http://dx.doi.org/10.1007/s00208-004-0618-y>. MR2139252.
- [Fra61] TH. FRANKEL, *Manifolds with positive curvature*, *Pacific J. Math.* **11** (1961), no. 1, 165–174. <http://dx.doi.org/10.2140/pjm.1961.11.165>. MR0123272.
- [FRW13] PH. FRANK, X. RONG, AND Y. WANG, *Fundamental groups of positively curved manifolds with symmetry*, *Math. Ann.* **355** (2013), no. 4, 1425–1441. <http://dx.doi.org/10.1007/s00208-012-0823-z>. MR3037020.
- [FZ11] L. A. FLORIT AND W. ZILLER, *Topological obstructions to fatness*, *Geom. Topol.* **15** (2011), no. 2, 891–925. <http://dx.doi.org/10.2140/gt.2011.15.891>. MR2821566.
- [GGS11] F. GALAZ-GARCIA AND C. SEARLE, *Low-dimensional manifolds with non-negative curvature and maximal symmetry rank*, *Proc. Amer. Math. Soc.* **139** (2011), no. 7, 2559–2564. <http://dx.doi.org/10.1090/S0002-9939-2010-10655-X>. MR2784821.
- [GGS12] F. GALAZ-GARCIA AND W. SPINDELER, *Nonnegatively curved fixed point homogeneous 5-manifolds*, *Ann. Global Anal. Geom.* **41** (2012), no. 2, 253–263. <http://dx.doi.org/10.1007/s10455-011-9282-0>. MR2876698.
- [GGS14] F. GALAZ-GARCIA AND C. SEARLE, *Nonnegatively curved 5-manifolds with almost maximal symmetry rank*, *Geom. Topol.* **18** (2014), no. 3, 1397–1435. <http://dx.doi.org/10.2140/gt.2014.18.1397>. MR3228455.
- [GK04] K. GROVE AND C. W. KIM, *Positively curved manifolds with low fixed point cohomogeneity*, *J. Differential Geom.* **67** (2004), no. 1, 1–33. MR2153480.
- [GM69] D. GROMOLL AND W. MEYER, *On complete open manifolds of positive curvature*, *Ann. of Math.* (2) **90** (1969), no. 1, 75–90. <http://dx.doi.org/10.2307/1970682>. MR0247590.
- [GoZ15] F. J. GOZZI, *Low dimensional polar actions*, *Geom. Dedicata* **175** (2015), 219–247. <http://dx.doi.org/10.1007/s10711-014-0037-5>. MR3323638.
- [Gro09] K. GROVE, *Developments around positive sectional curvature*, *Surveys in Differential Geometry*, Vol. XIII. Geometry, Analysis, and Algebraic Geometry: Forty Years of the Journal of Differential Geometry, *Surv. Differ. Geom.*, vol. 13, Int. Press, Somerville, MA, 2009, pp. 117–133. <http://dx.doi.org/10.4310/SDG.2008.v13.n1.a4>. MR2537084.

- [GS94] K. GROVE AND C. SEARLE, *Positively curved manifolds with maximal symmetry-rank*, J. Pure Appl. Algebra **91** (1994), no. 1–3, 137–142. [http://dx.doi.org/10.1016/0022-4049\(94\)90138-4](http://dx.doi.org/10.1016/0022-4049(94)90138-4). MR1255926.
- [GS97] ———, *Di erential topological restrictions curvature and symmetry*, J. Di erential Geom. **47** (1997), no. 3, 530–559. MR1617636.
- [GVZ11] K. GROVE, L. VERDIANI, AND W. ZILLER, *An exotic  $T_1S^4$  with positive curvature*, Geom. Funct. Anal. **21** (2011), no. 3, 499–524. <http://dx.doi.org/10.1007/s00039-011-0117-8>. MR2810857.
- [GW14] K. GROVE AND B. WILKING, *A knot characterization and 1-connected nonnegatively curved 4-manifolds with circle symmetry*, Geom. Topol. **18** (2014), no. 5, 3091–3110. <http://dx.doi.org/10.2140/gt.2014.18.3091>. MR3285230.
- [GWZ08] K. GROVE, B. WILKING, AND W. ZILLER, *Positively curved cohomogeneity one manifolds and 3-Sasakian geometry*, J. Di erential Geom. **78** (2008), no. 1, 33–111. MR2406265.
- [HK89] W. Y. HSIANG AND B. KLEINER, *On the topology of positively curved 4-manifolds with symmetry*, J. Di erential Geom. **29** (1989), no. 3, 615–621. MR992332.
- [HPW15] C. HE, P. PETERSEN, AND W. WYLIE, *Warped product Einstein metrics on homogeneous spaces and homogeneous Ricci solitons*, J. Reine Angew. Math. **707** (2015), 217–245. <http://dx.doi.org/10.1515/crelle-2013-0078>. MR3403459.
- [Ken] L. KENNARD, *Fundamental groups of manifolds with positive sectional curvature and torus symmetry*, J. Geom. Anal., to appear, available at <http://arxiv.org/abs/arXiv:1310.7251>.
- [Ken13] ———, *On the Hopf conjecture with symmetry*, Geom. Topol. **17** (2013), no. 1, 563–593. <http://dx.doi.org/10.2140/gt.2013.17.563>. MR3039770.
- [Ken14] ———, *Positively curved Riemannian metrics with logarithmic symmetry rank bounds*, Comment. Math. Helv. **89** (2014), no. 4, 937–962. <http://dx.doi.org/10.4171/CMH/340>. MR3284301.
- [Ker11] M. KERIN, *Some new examples with almost positive curvature*, Geom. Topol. **15** (2011), no. 1, 217–260. <http://dx.doi.org/10.2140/gt.2011.15.217>. MR2776844.
- [KM] A. V. KOLESNIKOV AND E. MILMAN, *Poincaré and Brunn-Minkowski inequalities on weighted Riemannian manifolds with boundary*, preprint, available at <http://arxiv.org/abs/arXiv:1310.2526>.
- [KT14] M. M. KERR AND K. TAPP, *A note on quasi-positive curvature conditions*, Di erential Geom. Appl. **34** (2014), 63–79. <http://dx.doi.org/10.1016/j.difgeo.2014.03.009>. MR3209537.
- [Lic70] A. LICHNEROWICZ, *Variétés riemanniennes à tenseur  $C$  non négatif*, C. R. Acad. Sci. Paris Sér. A-B **271** (1970), A650–A653 (French). MR0268812.
- [Lic72] ———, *Variétés kählériennes à première classe de Chern non négative et variétés riemanniennes à courbure de Ricci généralisée non négative*, J. Di erential Geometry **6** (1971/72), 47–94 (French). MR0300228.
- [Lot03] J. LOTT, *Some geometric properties of the Bakry-Émery-Ricci tensor*, Comment. Math. Helv. **78** (2003), no. 4, 865–883. <http://dx.doi.org/10.1007/s00014-003-0775-8>. MR2016700.
- [LV09] J. LOTT AND C. VILLANI, *Ricci curvature for metric-measure spaces via optimal transport*, Ann. of Math. (2) **169** (2009), no. 3, 903–991. <http://dx.doi.org/10.4007/annals.2009.169.903>. MR2480619.
- [Mor05] F. MORGAN, *Manifolds with density*, Notices Amer. Math. Soc. **52** (2005), no. 8, 853–858. MR2161354.
- [Mor06] ———, *Myers’ theorem with density*, Kodai Math. J. **29** (2006), no. 3, 455–461. <http://dx.doi.org/10.2996/kmj/1162478772>. MR2278776.
- [Mor09a] ———, *Geometric Measure Theory: A Beginner’s Guide*, 4th ed., Elsevier/Academic Press, Amsterdam, 2009. MR2455580.

- [Mor09b] ———, *Manifolds with density and Perelman's proof of the Poincaré conjecture*, Amer. Math. Monthly **116** (2009), no. 2, 134–142. <http://dx.doi.org/10.4169/193009709X469896>. MR2478057.
- [MW12] O. MUNTEANU AND J. WANG, *Analysis of weighted Laplacian and applications to Ricci solitons*, Comm. Anal. Geom. **20** (2012), no. 1, 55–94. <http://dx.doi.org/10.4310/CAG.2012.v20.n1.a3>. MR2903101.
- [Oht] S. OHTA, *(K,N)-convexity and the curvature-dimension condition for negative N*, J. Geom. Anal. **26** (2016), no. 3, 2067–2096. <http://dx.doi.org/10.1007/s12220-015-9619-1>. MR3511469.
- [Per] G. PERELMAN, *The entropy formula for the Ricci flow and its geometric applications*, preprint, available at <http://arxiv.org/abs/arXiv:math.DG/0211159>.
- [Pet06] P. PETERSEN, *Riemannian Geometry*, 2nd ed., Graduate Texts in Mathematics, vol. 171, Springer, New York, 2006. MR2243772.
- [Ron02] X. RONG, *Positively curved manifolds with almost maximal symmetry rank*, Geom. Dedicata **95** (2002), 157–182, Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part II (Haifa, 2000). <http://dx.doi.org/10.1023/A:1021242512463>. MR1950889.
- [Sim] F. SIMAS, *Nonnegatively curved five-manifolds with non-abelian symmetry*, Geom. Dedicata **181** (2016), 61–82, available at <http://arxiv.org/abs/arXiv:1212.5022>. <http://dx.doi.org/10.1007/s10711-015-0112-6>. MR3475738.
- [Spi14] W. L. SPINDELER,  *$S^1$ -actions on 4-manifolds and fixed point homogeneous manifolds of non-negative curvature*, Ph.D. thesis (2014), Westfälischen Wilhelms-Universität Münster.
- [Stu06a] K. TH. STURM, *On the geometry of metric measure spaces. I*, Acta Math. **196** (2006), no. 1, 65–131. <http://dx.doi.org/10.1007/s11511-006-0002-8>. MR2237206.
- [Stu06b] ———, *On the geometry of metric measure spaces. II*, Acta Math. **196** (2006), no. 1, 133–177. <http://dx.doi.org/10.1007/s11511-006-0003-7>. MR2237207.
- [SW09] H. SUN AND Y. WANG, *Positive sectional curvature, symmetry and Chern's conjecture*, Differential Geom. Appl. **27** (2009), no. 1, 129–136. <http://dx.doi.org/10.1016/j.difgeo.2008.10.004>. MR2488996.
- [Ver04] L. VERDIANI, *Cohomogeneity one manifolds of even dimension with strictly positive sectional curvature*, J. Differential Geom. **68** (2004), no. 1, 31–72. MR2152908.
- [vRS05] M. K. VON RENESSE AND K. TH. STURM, *Transport inequalities, gradient estimates, entropy, and Ricci curvature*, Comm. Pure Appl. Math. **58** (2005), no. 7, 923–940. <http://dx.doi.org/10.1002/cpa.20060>. MR2142879.
- [VZ14] L. VERDIANI AND W. ZILLER, *Concavity and rigidity in non-negative curvature*, J. Differential Geom. **97** (2014), no. 2, 349–375. MR3263509.
- [Wal72] N. R. WALLACH, *Compact homogeneous Riemannian manifolds with strictly positive curvature*, Ann. of Math. (2) **96** (1972), no. 2, 277–295. <http://dx.doi.org/10.2307/1970789>. MR0307122.
- [Wei] WEISSKOPF, N., *Positive curvature and the elliptic genus*, New York J. Math. **23**, 193–212. <http://nyjm.albany.edu/j/2017/23-11v.pdf>.
- [Wei68] A. WEINSTEIN, *A fixed point theorem for positively curved manifolds*, J. Math. Mech. **18** (1968/1969), no. 2, 149–153. <http://dx.doi.org/10.1512/iumj.1969.18.18016>. MR0227894.
- [Wei80] ———, *Fat bundles and symplectic manifolds*, Adv. in Math. **37** (1980), no. 3, 239–250. [http://dx.doi.org/10.1016/0001-8708\(80\)90035-3](http://dx.doi.org/10.1016/0001-8708(80)90035-3). MR591728.
- [Wie15] M. WIEMELER, *Torus manifolds and non-negative curvature*, J. Lond. Math. Soc. (2) **91** (2015), no. 3, 667–692. <http://dx.doi.org/10.1112/jlms/jdv008>. MR3355120.
- [Wil03] B. WILKING, *Torus actions on manifolds of positive sectional curvature*, Acta Math. **191** (2003), no. 2, 259–297. <http://dx.doi.org/10.1007/BF02392966>. MR2051400.
- [Wil06] ———, *Positively curved manifolds with symmetry*, Ann. of Math. (2) **163** (2006), no. 2, 607–668. <http://dx.doi.org/10.4007/annals.2006.163.607>. MR2199227.

- [Wil07] ———, *Nonnegatively and positively curved manifolds*, Surveys in Differential Geometry. Vol. XI, Int. Press, Somerville, MA, 2007, pp. 25–62. <http://dx.doi.org/10.4310/SDG.2006.v11.n1.a3>. MR2408263.
- [WW09] G. WEI AND W. WYLIE, *Comparison geometry for the Bakry-Emery Ricci tensor*, J. Differential Geom. **83** (2009), no. 2, 377–405. MR2577473.
- [Wy108] W. WYLIE, *Complete shrinking Ricci solitons have finite fundamental group*, Proc. Amer. Math. Soc. **136** (2008), no. 5, 1803–1806. <http://dx.doi.org/10.1090/S0002-9939-07-09174-5>. MR2373611.
- [Wy115] ———, *Sectional curvature for Riemannian manifolds with density*, Geom. Dedicata **178** (2015), 151–169. <http://dx.doi.org/10.1007/s10711-015-0050-3>. MR3397488.
- [Zil07] W. ZILLER, *Examples of Riemannian manifolds with non-negative sectional curvature*, Surveys in Differential Geometry. Vol. XI, Int. Press, Somerville, MA, 2007, pp. 63–102. <http://dx.doi.org/10.4310/SDG.2006.v11.n1.a4>. MR2408264.
- [Zil14] ———, *Riemannian manifolds with positive sectional curvature*, Geometry of Manifolds with Non-negative Sectional Curvature, Lecture Notes in Math., vol. 2110, Springer, 2014, pp. 1–19. [http://dx.doi.org/10.1007/978-3-319-06373-7\\_1](http://dx.doi.org/10.1007/978-3-319-06373-7_1). MR3329927.

LEE KENNARD:  
 Department of Mathematics  
 University of Oklahoma  
 Norman, OK 73019, USA  
 E-MAIL: [kennard@math.ou.edu](mailto:kennard@math.ou.edu)  
 URL: <http://www2.math.ou.edu/~kennard/>

WILLIAM WYLIE:  
 215 Carnegie Building  
 Department of Mathematics  
 Syracuse University  
 Syracuse, NY, 13244, USA  
 E-MAIL: [wwylie@syr.edu](mailto:wwylie@syr.edu)  
 URL: <https://wwylie.expressions.syr.edu>

KEY WORDS AND PHRASES: Positive sectional curvature, manifolds with density, symmetry.

2010 MATHEMATICS SUBJECT CLASSIFICATION: 53C20.

*Received: June 8, 2015.*